

# Roane County Office of Emergency Services

## EMS Division

### Protocol Guidelines



Update Effective February 1, 2016

# Roane County OES, EMS Division

---

FEB 2016

## TABLE OF CONTENTS

Administrative Notes.....	5
Clinical Notes.....	7
Clinical Notes - Airway.....	8
Clinical Notes - Cardio Vascular.....	8
Clinical Notes - IV.....	9
Definitions.....	10

## CARDIAC EMERGENCY

101	Automatic External Defibrillator (AED).....	11
102	Atrial Fibrillation and Flutter.....	12
	103	
	Bradycardia.....	13
104	Acute Coronary Syndrome/STEMI.....	15
105	Chest Pain / NON Cardiac.....	17
106	Pulseless Electrical Activity (P.E.A.).....	19
107	Premature Ventricular Contractions (PVC's).....	20
108	Supraventricular Tachycardia (SVT).....	21
109	Torsades de Pointes.....	23
110	Ventricular Asystole.....	25
111	Ventricular Fibrillation / Pulseless Ventricular Tachycardia.....	26
112	Ventricular Tachycardia with a Pulse .....	27
113	Post Resuscitation.....	29

## ENVIRONMENTAL EMERGENCY

201	Chemical Exposure .....	30
202	Drug Ingestion.....	31
203	Electrocution / Lightning Injuries.....	32
204	Hyperthermia.....	33
205	Hypothermia.....	35
206	Near Drowning.....	37
207	Nerve Agent Exposure .....	38
208	Poisonous Snake Bite.....	40
209	Radiation/Hazmat.....	41
210	Carbon Monoxide Exposure.....	42

## MEDICAL EMERGENCY

300	Medical Complaint Not Specified under other Protocols.....	43
301	Abdominal Pain (non-traumatic)/Nausea and Vomiting.....	44
302	Acute Pulmonary Edema / CHF.....	45

# Roane County OES, EMS Division

---

303	Anaphylactic Shock.....	46
304	Cerebrovascular Accident (CVA).....	47
	*Cincinnati Prehospital Stroke Scale.....	48
	*Thrombolytic Screening .....	49
305	Croup.....	50
306	Family Violence.....	51
307	Hyperglycemia Associated with Diabetes.....	52
309	Hypoglycemia.....	53
310	Medications at Schools.....	54
311	Non-Formulary Medications.....	55
312	Respiratory Distress (Asthma/COPD).....	56
313	Seizures.....	57
314	Sexual Assault.....	58
315	Sickle Cell Anemia.....	59
316	Unconscious / Unresponsive / Altered Mental Status.....	60
317	Syncope.....	61
318	Pain Management.....	62

## TRAUMA EMERGENCY

401	Air Ambulance Transport.....	63
402	Abdominal/Pelvic Trauma.....	64
403	Avulsed Teeth - Standing Order.....	66
404	Cardiogenic Shock.....	67
405	Eye Trauma.....	68
406	Hypovolemic Shock.....	69
407	Major Thermal Burn.....	71
408	Musculoskeletal Trauma.....	73
409	Multi-System Trauma.....	75
410	Neurogenic Shock.....	76
411	Septic Shock.....	77
412	Soft Tissue / Crush Injuries.....	78
413	Spinal Cord Injuries.....	80
414	Traumatic Cardiac Arrest.....	81
415	Tension Pneumothorax.....	82
416	Traumatic Amputation(s).....	84

## OBSTETRICAL EMERGENCIES

500	Obstetrical / Gynecological Complaints (Non-Delivery or Gynecological Only).....	85
501	Normal Delivery.....	86
	*APGAR scoring chart.....	87
502	Abruptio Placenta.....	89
503	Amniotic Sac Presentation.....	90
504	Breech or Limb Presentation.....	91
505	Meconium Stain.....	92
506	Placenta Previa.....	93
507	Prolapsed Umbilical Cord.....	94

# Roane County OES, EMS Division

---

508	Pre-eclampsia and Eclampsia.....	95
-----	----------------------------------	----

## MISCELLANEOUS EMERGENCY

601	Discontinuation / Withholding of Life Support.....	96
602	Field Determination of Death .....	97
603	Mandatory EKG.....	98
604	Patient Refusal or Declination of Care / Patient Non-Transport Situations.....	99
605	Physical Restraint.....	101
606	Physician On Scene.....	103
607	By-standers On Scene.....	104
608	Procedure for Deviation from Standing Orders.....	104
609	Spinal Immobilization.....	105
610	Stretcher Transport.....	110
611	Terminally Ill Patients .....	111
612	“Excited Delirium” / Taser Use.....	112
613	Neonatal Resuscitation.....	114

## PROCEDURES

Capnography.....	116
Chest Decompression.....	117
Continuous Positive Airway Pressure (CPAP).....	118
Delayed Off Loading of Stable Non-Emergent Patients in the ED .....	120
Endotracheal Tube Introducer (Bougie).....	121
External Transcutaneous Cardiac Pacing.....	122
EZ-IO.....	123
Intranasal Medication.....	126
Nu-Trake.....	129
RSI / DAI.....	130
Tourniquet.....	133

## REFERENCE

Trauma Score.....	134
Glasgow Coma Scale.....	135
Medical Abbreviations.....	136
Drug Infusion Dosage Guidelines.....	137
Pediatric Trauma Score.....	138

## Introduction

These Protocol guidelines are provided by State of Tennessee Office of Emergency Medical Services and are designed to be used as written or as a guideline for Emergency Medical Directors of Licensed Emergency Medical Services in Tennessee. Protocols provide direction for Emergency Medical Services Personnel to render appropriate care for the sick and injured of all ages. It is recommended that services require EMS Personnel to familiarize themselves with the service approved Protocols and show successful completion by written documentation of competency in the Service Protocols to the Service Medical Director.

# Roane County OES, EMS Division

---

## Administrative Notes:

1. The EMT and Advanced EMT (AEMT) will assist ALS personnel as requested and/or needed.
2. The Emergency Medical Responder will function under the current guidelines as stated in the AHA-BLS Healthcare Provider text. Shall also be responsible for other duties as assigned within their Scope of Practice by the AEMT or the Paramedic.
3. Providers currently licensed as AEMT will continue to function at their current scope of practice until the appropriate “bridge” certification has been obtained through a state accredited program.
4. The Paramedic will be in charge and will be responsible for all of the actions and or activities as it relates to the Emergency Unit. On the scene of an emergency, the Paramedic will be responsible for patient care. The EMT or AEMT will act within their scope of practice to any request for patient care or maintenance of the unit as directed by the Paramedic. Patient care is limited to acts within their scope of practice as defined by these SOPs. The EMS Personnel are responsible for reviewing all documentation and signing in the required manner
5. It is the responsibility of the most qualified Paramedic caring for the patient to ensure transmission of all aspects of the patient assessment and care to the responding Emergency Unit or Medical Control.
6. When reporting a disposition to Medical Control or the responding unit, provide the following minimum information:
  - A. Patient’s age and chief complaint
  - B. Is the patient stable or unstable, including complete V/S and LOC
  - C. Interventions performed
  - D. Provide other information as requested.
7. For each and every call, the first directives are scene safety and body substance isolation precautions.
8. For any drug administration of procedures outside these Guidelines, the EMS Provider must receive authorization from Medical Control. Paramedics en-route to the scene are not authorized to issue orders.
9. The minimal equipment required for all patient calls:
  - A. When the patient is in close proximity to the unit or Emergency Medical Responder: jump bag, cardiac monitor, and oxygen or other equipment as may be indicated by the nature of the call
  - (Cont. on next page)
  - B. When the patient is not in close proximity of the unit or Emergency Medical Responder: the above equipment, stretcher and any other equipment that may be needed as dictated by the nature of the call.
10. The senior Paramedic has the ultimate responsibility to ensure that all records and reports are properly completed. The patient care report should accurately reflect the clinical activities undertaken. If there is a patient refusal, declination, or dismissal of service at the scene of the incident, the incident report should reflect the details as well as the party or parties responsible for the request to terminate any and all evaluations and treatment.

## Roane County OES, EMS Division

---

11. Although the Guidelines have a numerical order, it may be necessary to change the sequence order or even omit a procedure due to patient condition, the availability of assistance, or equipment. Document your reason for any deviations from protocol.
12. EMTs and AEMTS are expected to perform their duties in accordance with local, state and federal guidelines in accordance with the State of Tennessee statutes and rules of Tennessee Emergency Services. The Paramedic will work within their scope of practice dependent on available equipment.
13. Each patient care contact will be recorded on the EMS patient care report as completely and accurately as practical and per agency guidelines. A complete copy of the patient out-of-hospital evaluation(s) and treatment(s) will be given to the emergency department personnel or staff prior to departing from the health care facility. This will ensure proper documentation of the continuity of care.
14. In potential crime scenes, any movement of the body, clothing, or immediate surroundings should be documented and the on scene law enforcement officer should be notified of such.
15. All patients should be transported to the most appropriate facility according to the patient or family request or the facility that has the level of care commensurate with the patient's condition. Certain medical emergencies may require transport to a facility with specialized capability.
16. Paramedics may transport the patient in a non-emergency status to the hospital. This should be based on the signs and symptoms of the patient, mechanism of injury or nature of illness.
17. The following refusal situations should be evaluated by a paramedic.
  - A. Hypoglycemic patients who have responded to treatment
  - B. Any patient refusing transport who has a potentially serious illness or injury
  - C. Patients age less than 4 years or greater than 70 years
  - D. Chest pain any age or cause
  - E. Drug overdose / intoxicated patients
  - F. Potentially head injured patients
18. The use of a length based assessment tape is **required** for all pediatric patients as a guide for medications and equipment sizes. The tape will be utilized on all pediatric patients below the age of 8 years and appropriate for their weight. Any child that is small in stature for their age, you should consider utilizing the length based tape for compiling a complete accurate assessment of the patient. This information will be passed along to the receiving facility and documented in the PCR.

# Roane County OES, EMS Division

---

## Clinical Notes:

1. A complete patient assessment, vital signs, treatments and continued patient evaluation are to be initiated immediately upon contact with patient and continued until patient care is transferred to a Higher Medical Authority. Refer to Patient Assessment Flow Chart.
2. The on-going assessment times are considered:

<u>High Priority</u>	<u>Low Priority</u>
Every 3 - 5 minutes	Every 5 - 15 minutes
3. EMTs may utilize the following medications: Aspirin, Nitroglycerine, Epinephrine (for Anaphylactic reaction), and assist patient with their own Albuterol or MDI. AEMTs may administer Albuterol, MDI, and Dextrose for hypoglycemia as well as other medications within their scope of practice. Use Nitroglycerine with caution in patients taking erectile dysfunction medications as profound hypotension may occur.
4. If a glucometer reading of greater than 40 mg/dL and patient is asymptomatic, start an INT and administer oral glucose. If a glucometer reading is less than 80 mg/dL and patient is symptomatic, start an IV NS and administer 12.5 - 25 gms of Dextrose. Reassess patient every 5 min, repeat PRN  
**Note:** Any administration of Dextrose must be done through an IV line, not INTs.  
Normal blood sugar values are 80 - 120 mg/dL.
5. Blood Glucose and Stroke Screening will be performed on all patients with altered mental status. Glucose should be titrated slowly in order to restore normal levels while avoiding large changes in serum glucose levels. Be aware that elevated glucose levels are detrimental in conditions such as stroke.
6. Supportive care indicates any emotional and/or physical care including oxygen therapy, repositioning patient, comfort measures and patient family education.
7. Upon arrival at the receiving hospital, all treatment(s) initiated in the field will be continued until hospital personnel have assumed patient care.
8. The initial blood pressure **MUST** be taken manually. If subsequent blood pressures taken by machine vary more than 15 points diastolic, then a manual blood pressure will verify the machine reading.
9. EMTs may obtain and transmit EKG monitoring tracings and 12 Lead EKGs in the presence of the treating Paramedic. Paramedics **ONLY** may interpret and make treatment and destination decisions based on the 12 lead EKG.
10. Indications for football helmet removal:
  - When a patient is wearing a helmet and not shoulder pads
  - In the presence of head and/or facial trauma
  - Patients requiring advanced airway management when removal of the facemask is not sufficient
  - When the helmet is loose on the patient's head
  - In the presence of cardiopulmonary arrest. (The shoulder pads must also be removed)

## Roane County OES, EMS Division

---

When helmet and shoulder pads are both on the spine is kept in neutral alignment. If the patient is wearing only a helmet or shoulder pads, neutral alignment must be maintained. Either remove the other piece of equipment or pad under the missing piece. *All other helmets must be removed in order to maintain spinal alignment.*

### Clinical Notes - Airway:

1. All EMTs have standing orders for insertion of an approved airway device for patients meeting the indications
2. Airway maintenance appropriate for the patient's condition includes any airway maneuver, adjunct, or insertions of tubes that provide a patent airway.
3. Pulse Oximetry should be utilized for all patients complaining of respiratory distress or chest pain (regardless of source).
4. Esophageal Intubation detectors and End Tidal CO<sub>2</sub> or waveform capnography (preferred) are **MANDATORY** for all intubations. Reliability may be limited in patients less than 20kg. Use other methods to assist in confirmation.
5. Use of cervical collars post intubation (BIAD or ET) is recommended to reduce the chance of accidental extubation. This is in addition to the tube securing devices currently in use.

### Clinical Notes - Cardio-Vascular

1. In the adult cardiac arrest:
  - A. CPR is most effective when done continuously, with minimum interruption. Maintain rate of 110 BPM, a depth of 2 inches and a compression fraction of >80%.
  - B. Initiate compressions first, manage airway after effective compressions for two minutes
  - C. All IV/IO drugs given are to be followed by a 10 cc NS bolus
  - D. Elevate the extremity after bolus when given IV
  - E. Consider blind airway devices (King) whenever intubation takes longer than 30 seconds
  - F. Apply NC Oxygen 2 - 4 L during initial CPR
  - G. Consider use of mechanical CPR device if available
2. Treat the patient not the monitor
3. Defibrillation and Synchronized Cardioversion joules are based on the use of the current biphasic monitor.
4. If a change in cardiac rhythm occurs, provide all treatment and intervention as appropriate for the new rhythm.
5. In the case of cardiac arrest where venous access is not readily available, paramedics may use EZ-IO as initial access. Humeral access is preferred in medical conditions.
6. Place all cardiac patients on a heart monitor and appropriate personnel obtain a 12 lead EKG if any concern for an acute cardiac disturbance.

## Roane County OES, EMS Division

---

### Clinical Notes - IV

1. AEMTs and Paramedics have standing orders for precautionary IV and INTs. AEMTs have a standing order for the insertion of an IV or INT under the following guidelines:
  - A. The patient must have some indication that they are unstable (see definitions)
  - B. Limited to two attempts in one arm only. (Cannulation of legs or neck is not allowed.)
  - C. Drug administration will be followed by a minimum of 10cc of fluid to flush the catheter.
  - D. Blood Glucose will be performed for all patients with altered mental status
  - E. IVs should not be attempted in an injured extremity
  - F. TKO (To Keep Open) indicates a flow rate of approximately 50 cc/hr G. IVs will not be started in arms with shunts
    - H. IVs appropriate for patients condition:
      - I. if patient is hypotensive, give a bolus of fluid
      - II. if patient's BP is normal run IV TKO or convert to saline lock (INT). III. A bolus of fluid is 20 cc/kg for all patients.
2. For external Jugular IVs attempted by paramedics, IV catheters should be 18 gauge or smaller diameter based on the patient.
3. Paramedics, when properly equipped and trained, may utilize indwelling access ports such as Port-A-Cath in an **EMERGENCY ONLY**. This procedure should be done with a Huber needle utilizing sterile technique.

# Roane County OES, EMS Division

---

## Definitions

**Standing Order** - This skill or treatment **may** be initiated prior to contact with Medical Control.

**Protocol** - A suggested list of treatment options **requiring** you to contact Medical Control **prior** to initiation

**Medical Director** - the physician that has the ultimate responsibility for the patient care aspects of the EMS System

**Unstable (symptomatic)** - indicates that one or more of the following are present:

- a. Chest pain / Dyspnea
- b. Hypotension (systolic B/P less than 90 mmHg in a 70 kg patient or greater)
- c. Signs and symptoms of congestive heart failure or pulmonary edema
- d. Signs and symptoms of a myocardial infarction
- e. Signs and symptoms of inadequate perfusion
- f. Altered level of consciousness

**Stable (asymptomatic)** - Indicates that the patient has no or very mild signs and symptoms associated with the current history of illness or trauma.

**Emergency Medical Responder** - Personnel licensed by the Tennessee Department of Health, Office of EMS and authorized by the service Medical Director to perform lifesaving interventions while awaiting additional EMS response. May also assist higher level personnel at scene and during transport under medical direction and within their scope of practice.

**EMT** - Personnel licensed by the Tennessee Department of Health, Office of EMS and authorized by the Medical Director to provide basic emergency care according to the Standard of Care and these Guidelines.

**AEMT** - Personnel licensed by the Tennessee Department of Health, Office of EMS and authorized by the Medical Director to provide limited advanced emergency care according to the Standard of Care and Standing Orders and Protocols.

**Paramedic** - Personnel licensed by the Tennessee Department of Health, Office of EMS and authorized by the Medical Director to provide basic and advanced emergency patient care according to the standard of care and these guidelines Orders and Protocols

**Transfer of Care** - Properly maintaining the continuity of care through appropriate verbal and/or written communication of patient care aspects to an equal or higher appropriate medical authority.

**Higher Medical Authority** - Any medical personnel that possesses a current medical license or certificate recognized by the State of Tennessee with a higher level of medical training than the one possessed by EMS Personnel. (MD)

**Medical Control (transport)** - The instructions and advice provided by a physician, and the orders by a physician that define the treatment of the patient. To access Medical Control, contact the Emergency Department physician on duty of the patient's first choice.

# Roane County OES, EMS Division

## CARDIAC EMERGENCY

\*\*\*\*All EMRs, EMTs, AEMTs, and Paramedics are expected to perform their duties in accordance with local, state and federal guidelines.

### 101 Automatic External Defibrillator (AED)

#### Assessment

Patient in Cardiopulmonary Arrest  
Basic Life Support in progress AED  
in use

#### EMR and EMT

1. If AED available, apply to patient and follow prompts
2. 100% oxygen and airway maintenance appropriate to patient's condition. All CPR rates of compression are 100 per minute for all ages. Ventilation rates are 8 - 10/min with no pause in CPR for ventilation.
3. Continue CPR according to current AHA - Healthcare Provider Guidelines, specific for patient's age.
4. If AED is in use (defibrillating) prior to arrival, allow shocks to be completed, and then elevate pulse.
  - A. If no pulse, continue to provide CPR and basic life support.
  - B. If a pulse is present, evaluate respirations and provide supportive care appropriate for the patient's condition.
5. **EMR and EMT STOP**

#### AEMT

6. IV NS Bolus (20 cc/kg), then TKO
7. **AEMT STOP**

#### PARAMEDIC

8. Monitor patient and treat per SOP specific for the arrhythmia

#### Notes:

1. AED is relatively **contraindicated** in the following situations:
  - A. If the victim is in standing water, remove the victim from the water, and ensure that chest and surrounding area is dry.
  - B. Trauma Cardiac Arrest
2. Victims with implanted pacemakers, place pads 1 inch from device.
  - C. If ICD/AICD is delivering shock to the patient allow 30 to 60 seconds for the ICD/AICD to complete the treatment cycle before using the AED.
3. Transdermal medication patch at site of the AED pads:
  - D. If a medication patch is in the location for an AED pad, remove the medication patch and wipe the area clean before attaching the AED electrode pad.

## 102 Atrial Fibrillation and Flutter

### Assessment

Paroxysmal Atrial Tachycardia  
Atrial flutter **new onset**  
Atrial fibrillation **new onset**  
Symptomatic patient  
Dyspnea  
Chest pain  
Radiating pain  
Altered mental status  
Hypotension (systolic BP <90 mmHg) Diaphoresis

### EMR

1. Oxygen 100% and airway maintenance appropriate for the patient's condition
2. Supportive care
3. **EMR STOP**

### EMT

4. Pulse Oximetry
5. **EMT STOP**

### AEMT

6. Glucose check
7. IV NS TKO or INT
8. Titrate **Dextrose** 50% PRN slowly until normal levels achieved, if hypoglycemic
9. **AEMT STOP**

### PARAMEDIC

10. 12 Lead EKG
11. Valsalva maneuver
12. If perfusion is adequate, patient is stable, and HR is >150- No additional treatment required.
13. If perfusion is inadequate with HR > 150, administer **Diltiazem** 0.25 mg/kg IVP over 2 mins, may repeat after 10 min. @ 0.35 mg/kg IV Bolus.
14. If patient is unstable consider synchronized cardioversion\*\*\*:
  - A. Atrial flutter @ 30 joules (*pediatrics 0.5 j/kg then 1 j/kg*)
  - B. Atrial fib @ 50 joules (*pediatrics 0.5 j/kg then 1 j/kg*)
  - C. Pre-medicate with **Versed** 2-5 mg IVP (*pediatrics 0.1 mg/kg*) and/or **Morphine** 2-5 mg IV (*0.1 mg/kg*)

\*\*\*Immediate Synchronized Cardioversion (50, 75, 100, 120 150, 200 joules) (*pediatrics 0.5 j/kg then 1 j/kg*) is recommended when there is an unstable rhythm with serious signs and symptoms (CP, SOB, Decreased LOC Low Blood Pressure)

# Roane County OES, EMS Division

## CARDIAC EMERGENCY

### 103 Bradycardia

#### Assessment

Heart rate less than 60 beats per minute and symptomatic  
Decreased / altered LOC  
Chest pain / discomfort  
CHF / pulmonary edema  
Head Trauma  
Elevated Intracranial Pressure  
Dyspnea  
Hypothermia  
Hypoglycemia  
Drug overdose  
Signs of decreased perfusion  
Rhythm may be sinus bradycardia, junctional, or heart block Heart rates <80/min for infant or <60/min for child

#### EMR

1. Oxygen and airway maintenance appropriate to patient's condition. (High Flow O<sub>2</sub>.)
2. Supportive care
3. **EMR STOP**

#### EMT

4. Pulse Oximetry
5. **EMT STOP**

#### AEMT

6. Glucose Check
7. INT or IV NS TKO
8. Titrate **Dextrose** 50% PRN slowly until normal levels achieved, if hypoglycemic.
9. **AEMT STOP**

#### PARAMEDIC

10. 12 Lead EKG
11. If patient is asymptomatic and heart rate is less than 60 beats per minute, observe
12. If PVCs are present with bradycardia **DO NOT** administer an antiarrhythmic
13. Adults - (**SYMPTOMATIC**)
  - A. If systolic BP <90 mmHG and heart rate <60/min
    - I. If 2<sup>nd</sup> and 3<sup>rd</sup> degree blocks are present apply transcutaneous pacer pads per protocol.
    - II. Consider **Atropine** 0.5 mg (up to max 3 mg for adults)

(Continued on next page)

## Roane County OES, EMS Division

---

- B. If systolic BP < 90 mmHG and heart rate < 60/min continues
  - I. Notify Medical Control and begin cardiac pacing per protocol.
  - II. Consider Dopamine 2 - 20 mcg/kg/min as a continuous IV infusion.
- 14. **Pediatric -**
  - A. **Heart rates < 60/min with poor perfusion, respiratory distress or hypotension**
  - B. **Start chest compression, IV/IO**
    - I. **Epinephrine 1:10,000 0.01 mg/kg IV/IO q 4 min**
    - II. **Contact Medical Control**
      - a. **Consider external cardiac pacing**
      - b. **Consider Dopamine 2 - 20 mcg/kg/min as a continuous IV infusion to increase heart rate**

# Roane County OES, EMS Division

## CARDIAC EMERGENCY

### 104 Acute Coronary Syndrome/STEMI

#### Assessment

Determine quality, duration and radiation of pain Substernal Oppressive Chest Pain (crushing or squeezing) Nausea and/or vomiting Shortness of breath Cool, clammy skin Palpitations Anxiety or restlessness Abnormal pulse rate or rhythm History of Coronary Artery Disease or AMI Currently taking cardiac medications JVD Distal pulse for equality/strength to assess for Aneurysm Diaphoresis, pallor, cyanosis Breath sounds - congestion, rales, wheezing Motor deficits	<p><b>P</b> - Placement of pain/discomfort (anything that increases discomfort) <b>Q</b> - Quality of pain <b>R</b> - Radiation of pain <b>S</b> - Severity of pain/discomfort (scale of 1 - 10) <b>T</b> - Time of pain/discomfort onset, type of pain</p> <p>The elderly, women, and/or diabetic patients may complain of nausea, weakness, shortness of breath or other vague symptoms. Screen all such patients for possible silent MI</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

#### EMR

1. Oxygen at 2 - 6 LPM BNC and airway maintenance appropriate to patient's condition. If the patient is in severe respiratory distress consider Oxygen 100% 12 - 15 LPM NRB
2. Supportive Care
3. **EMR STOP**

#### EMT

4. Pulse oximetry, provide O<sub>2</sub> sufficient to keep SATs >94%
5. Give 324 mg of aspirin (chewable non enteric coated) if patient has no contraindications or has not already self-dosed. May assist with patient's NTG SL.
6. **EMT STOP**

#### AEMT

7. Glucose Check
8. INT or IV Normal Saline TKO
9. Titrate **Dextrose** 50% PRN slowly until normal levels achieved, if hypoglycemic
10. If systolic BP is >110 and the patient is symptomatic, assist patient with 1 NTG tablet or spray sublingually and reassess every 5 minutes. (Refer to the medication assist procedure.) Maximum of 3 doses.
11. Contact Medical Control to request orders for additional Nitroglycerin in excess of three doses.

## Roane County OES, EMS Division

---

**NOTE:** The maximum dosage of Nitroglycerin is three. The total dosage is the total doses the patient has taken on their own combined with your subsequent dosages. Use with caution in patients taking erectile dysfunction medications as this may cause profound hypotension

### 12. AEMT STOP

### PARAMEDIC

13. Cardiac monitor, obtain 12 lead
14. Patients with positive AMI should be transported to an appropriate cardiac facility as soon as possible. Treat arrhythmia appropriately. Call a **Code STEMI**.
15. **Aspirin** (non enteric coated), 324 mg chewed then swallowed if not self-dosed within last 24 hr
16. **Nitroglycerin** - If patient is not Hypotensive (BP <100 mmHg) administer one **Nitroglycerin**. Repeat **Nitroglycerine** once every 5 minutes after initial administration. \*\*\*Continue nitrate therapy until pain relieved or systolic BP<100 mmHg.
17. Systolic BP is <100 mmHg give 250 ml NS bolus (assess for signs of pulmonary congestion)
18. If chest pain/discomfort continues
  - A. Continue Nitrate therapy
  - B. Complete thrombolytic screening
  - C. If chest pain greater than 7 on scale of 1-10, consider **Morphine** 2-5mg until pain is tolerated by patient
  - D. Contact Medical Control
  - E. Transport

### EMS Cardiac “Medical Miranda”

“You are having a heart attack which is best cared for at a hospital that can treat you by opening up a blocked artery with a catheter. We strongly recommend taking you to one of those hospitals. Can we take you to one of those hospitals that provides these services?”

Give them choices based on normal referral patterns of the patient, family and closest PCI hospital. If yes, take them to the one they have chosen.

**If no, tell them you will take them to their chosen hospital but their care may be delayed due to their choice. If they change their mind, change destination hospital.**

# Roane County OES, EMS Division

## CARDIAC EMERGENCY

### 105 Chest Pain / NON Cardiac

#### Assessment

Determine quality, duration and radiation of pain Atypical Chest Pain NO Nausea and/or Vomiting NO Shortness of breath NO Cool, clammy skin History of chest injury, persistent cough NO History of Coronary Artery Disease or AMI NOT currently taking cardiac medications Distal pulse for equality/strength to assess for aneurysm NO Diaphoresis, pallor, cyanosis Normal Breath sounds	<b>P</b> - Placement of pain/discomfort (anything increase discomfort) <b>Q</b> - Quality of pain <b>R</b> - Radiation of pain <b>S</b> - Severity of pain/discomfort (scale of 1 - 10) <b>T</b> - Time of pain/discomfort onset, type of pain  The elderly, women, and/or diabetic patients may complain of nausea, weakness, shortness of breath or other vague symptoms. Screen all such patients for possible silent MI
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

#### EMR

1. Oxygen at 2 - 6 Lpm NC and airway maintenance appropriate to patient's condition. If the patient is in severe respiratory distress, consider Oxygen 100% 12 - 15 Lpm NRB
2. Supportive Care
3. **EMR STOP**

#### EMT

4. Pulse oximetry
5. Give **Aspirin** (325mg of chewable non-enteric coated if patient has not selfadministered in the last 24 hours.)
6. **EMT STOP**

#### AEMT

7. Glucose check
8. INT or IV Normal Saline TKO
9. Titrate **Dextrose** 50% PRN slowly until normal levels achieved, if hypoglycemic.
10. If systolic BP is >110 and the patient is symptomatic, **assist** patient with 1 **NTG** tablet or spray sublingually and reassess every 5 minutes. (Refer to the medication assist procedure.) Maximum of three doses.
11. Contact Medical Control to request orders for additional Nitroglycerin in excess of three doses  
**NOTE:** The maximum dosage of Nitroglycerin is three sublingual administrations, whether before or after your arrival. Use with caution in patients taking erectile dysfunction medications. Profound hypotension may occur.
12. **AEMT STOP**

(Paramedic cont. on next page)

## Roane County OES, EMS Division

---

### PARAMEDIC

13. EKG monitor, obtain 12 lead (transmit)
14. If no effect from max dose of nitro consider:
  - A. If chest pain greater the 7 on scale of 1-10, consider **Morphine** 2-5mg until pain is tolerated by patient
16. Contact Medical Control
17. Transport

**CAUTION:** Patients with true cardiac disease may have subtle, atypical symptoms. Always err on the side of patient safety.

# Roane County OES, EMS Division

## CARDIAC EMERGENCY

### 106 Pulseless Electrical Activity (P.E.A.)

#### Assessment

Presence of electrical cardiac rhythm without palpable pulse  
Confirm rhythm electrodes in two leads

#### EMR and EMT

1. Utilize AED if available
2. Oxygen 100% and airway maintenance appropriate to the patient's condition
3. CPR as indicated
4. **EMR and EMT STOP**

#### AEMT

5. Glucose Check if time allows
6. IV NS, bolus of fluid (20 cc/kg)
7. Titrate **Dextrose 50%** PRN slowly until normal levels achieved, if hypotensive
8. **AEMT STOP**

#### PARAMEDIC

9. EKG Monitor, 12 Lead EKG
10. **Epinephrine (1:10,000) 1mg IVP/IO (peds - Epinephrine (1:10,000) 0.01 mg/kg IV/IO q 4 min)**
11. Search for underlying cause of arrest and provide the related therapy:
  - A. Hypoxia - ensure adequate ventilation
  - B. Hypovolemia - fluid administration/fluid challenge (adult 20 cc/kg, peds 20 cc/kg bolus)
  - C. Cardiac tamponade (adult up to 2 liter bolus. peds 20 cc/kg bolus)
  - D. Tension pneumothorax - needle decompression
  - E. **KNOWN** hyperkalemia or tricyclic antidepressant overdose - **Sodium Bicarbonate** 1 mEq/kg, may repeat @ 0.5 mEq/kg q 10 min (peds 1 mEq/kg may repeat at 0.5 mEq/kg q 10 min) and **CaCl** 500mg IVP (peds 20mg/kg)
  - F. Acidosis in prolonged arrest: consider Sodium Bicarbonate 1 - 1.5 mEq/kg IV
  - G. Drug Overdose: **Narcan** 0.4 mg IV/IO/IM/IN titrated to adequate ventilation. (peds 0.1 mg/kg IV/IO/IM/IN titrated to adequate ventilation). Max dose up to 2mg.
  - H. Hypothermia: Initiate patient rewarming, avoid chest compressions if spontaneous circulation.
12. Consider External Cardiac Pacing per Protocol
13. PEA continues: Continue CPR, transport to appropriate facility.

# Roane County OES, EMS Division

## CARDIAC EMERGENCY

### 107 Premature Ventricular Contractions (PVC)

#### Assessment

Any PVC in acute MI setting with associated chest pain  
Multi-focal PVCs  
Unifocal and >15/min  
Salvos/couplets/runs of V-Tach (three or more PVCs in a row) and symptomatic  
PVCs occurring near the “T-wave”

#### EMR

1. Oxygen 100% and airway maintenance appropriate for the patient’s
2. condition Supportive care
3. **EMR STOP**

#### EMT

4. Pulse oximetry
5. **EMT STOP**

#### AEMT

6. Glucose check
7. INT or IV NS TKO
8. Titrate Dextrose 50% PRN slowly until normal levels achieved, if hypotensive.
9. **AEMT STOP**

#### PARAMEDIC

10. EKG monitor, 12 lead EKG
11. If PVCs are present with heart rate >60/min
  - a. Treat underlying causes first
  - b. If still symptomatic then, **Amiodarone** 150 mg over 10 minutes. May repeat for a total of three doses if needed.

# Roane County OES, EMS Division

## CARDIAC EMERGENCY

### 108 Supraventricular Tachycardia (SVT)

#### Assessment

Adult patients with heart rates in excess of 160 bpm (*peds rate >220 bpm*) (QRS width <12 sec [3 small blocks])  
Patients may exhibit symptoms of dyspnea, chest pain, radiating pain, altered mental status, hypotension (Systolic BP >90 mm/Hg)

#### EMR

1. Oxygen 100% and airway maintenance appropriate for the patient's condition.
2. Supportive care
3. **EMR STOP**

#### EMT

4. Pulse oximetry
5. **EMT STOP**

#### AEMT

6. Glucose check
7. INT or IV, NS TKO
8. Titrate **Dextrose** 50% PRN slowly until normal levels achieved, if hypoglycemic
9. **AEMT STOP**

#### PARAMEDIC

10. EKG Monitor, 12 Lead EKG
11. Valsalva maneuver for 10 seconds
12. **Adenosine** 12 mg rapid IVP (*peds 0.2 mg/kg IVP 12 mg max*). If no conversion, repeat 12 mg dose. (Flush with 10cc NS after each dose)
  - A. If rhythm does not convert to <150/min, or if patient is unstable or significantly symptomatic prepare for **synchronized cardioversion** Sedate as necessary:
    - i. **Versed** 2-5 mg IVP (*peds 0.1 mg/kg IVP max dose 5mg*)
  - B. If rhythm converts to rate <150/min: reassess for changes, maintain systolic BP >90 mmHg, transport, and contact Medical Control
- \*\*\*NOTE: Due to increased sensitivity to drug effects in heart transplant patients and those on Tegretol, give ½ the normal dose of Adenosine.
13. If SVT continues refractory to adenosine, and identified to be A-flutter or A-fib, administer **Diltiazem** 0.25 mg/kg IVP over 2 mins, may repeat after 10 min. @ 0.35 mg/kg IVP if time allows.

#### \*\*\*NOTES:

1. Adenosine is administered through large bore IV in the Antecubital Fossa
2. Other vagal maneuvers may include asking the patient to hold their breath, trendelenburg position.

## Roane County OES, EMS Division

---

### CARDIAC EMERGENCY

3. **Unstable SVT** may utilize synchronized cardioversion immediately in unstable patients prior to IV access. Assess the situation and make a good decision. Cardioversion hurts!
4. Significant symptoms include diaphoresis, hypotension, poor color or perfusion, mental status changes, chest pain >7/10

# Roane County OES, EMS Division

## CARDIAC EMERGENCY

### 109 Torsades de Pointes

#### Assessment

Decreased / altered LOC  
Dyspnea  
Chest Pain / discomfort, suspected AMI  
Hypotension (systolic BP <90 mmHg)  
CHF / Pulmonary edema  
Heart rate >160/min with QRS >.12 sec (3 small blocks, wide complex) and twisting of points

#### EMR

1. Oxygen 100% and airway maintenance appropriate for the patient's condition, pulse oximetry.
2. Supportive Care
3. **EMR STOP**

#### EMT

4. Pulse oximetry
5. **EMT STOP**

#### AEMT

6. Glucose check
7. INT or IV NS TKO
8. Titrate Dextrose 50% PRN slowly until normal levels achieved, if hypoglycemic.
9. **AEMT STOP**

#### PARAMEDIC

10. EKG Monitor, 12 Lead EKG
11. Systolic BP
  - A. If <90 mmHg - unstable/symptomatic:
    - I. Prepare for cardioversion at 100j, escalate as needed.
    - II. Sedation as necessary:  
Versed 2-5mg IV (peds 0.1mg/kg) and/or Pain Medications per the Chart below
      - a. If rate <160/min - monitor for changes, transport, Magnesium Sulfate 1 - 2 g IVP over 2 minutes
      - b. If rate >160/min - contact Medical Control, consider Amiodarone 150 - 300 mg IV/IO (peds 5 mg/kg), transport.
  - B. If >90 mmHg -stable/asymptomatic:
    - I. Magnesium Sulfate 1-2 gram IVP over 2 min
      - a. If rate <160/min - monitor for changes, Magnesium Sulfate may repeat 1-2 gram IVP over 2 minutes, transport

## Roane County OES, EMS Division

---

### CARDIAC EMERGENCY

- b. If rate >160/min - contact Medical Control, consider Amiodarone 150-300 mg IV/IO (*peds 5 mg/kg*), maintain systolic BP >90 mm/Hg, transport

# Roane County OES, EMS Division

## CARDIAC EMERGENCY

### 110 Ventricular Asystole

#### Assessment

No pulse or respirations  
Confirm cardiac rhythm with electrodes  
Record in two leads to confirm Asystole and to rule out fine V-Fib.

#### EMR and EMT

1. AED
2. CPR appropriate for patient age
3. Oxygen 100% and airway maintenance appropriate to patient's condition
4. **EMR and EMT STOP**

#### AEMT

5. Glucose check
6. IV NS bolus (20 cc/kg bolus fluids)
7. Titrate **Dextrose** 50% PRN slowly until normal levels achieved, if hypoglycemic
8. **AEMT STOP**

#### PARAMEDIC

9. Epinephrine 1:10,000 1 mg IO/IVP every 3-5 minutes (*peds Epinephrine 1:10,000 0.01 mg/kg IV/IO q 5 min*)
10. For prolonged resuscitation, consider: Sodium Bicarbonate 1 mEq/kg IV/IO followed by 0.5 mEq/kg q 10 min (*peds 1 mEq/kg may repeat at 0.5 mEq/kg q 10 min*).
11. Consider:
  - A. Magnesium Sulfate 1-2 gm Slow IV push over two minutes (*No peds dosing*)
  - B. Defibrillation for possible fine ventricular fibrillation masquerading as asystole
  - C. Consider external pacing under the following circumstances:
    - I. If cardiopulmonary arrest was witnessed by an experienced provider, and the patient is in asystole,
    - II. Prompt application of the transcutaneous cardiac pacemaker is appropriate prior to the administration of Epinephrine when a patient converts to asystole as a primary rhythm during EKG monitoring.
  - D. CaCl if arrest secondary to renal failure, or history of hemodialysis, adult 500 mg IV (*peds 20 mg/kg*)
  - E. Consider discontinuing efforts if criteria are met under Discontinuation/ Withholding of Life Support standing order.

#### Reversible Causes

Hypovolemia	Tamponade(Cardiac)
Hypoxia	Tension Pneumothorax
Hydrogen ion (acidosis)	Thrombosis(Heart)
Hyperkalemia/Hypokalemia	Thrombosis (Lungs)
Hypothermia	Tablets(Overdose)

# Roane County OES, EMS Division

## CARDIAC EMERGENCY

### 111 Ventricular Fibrillation / Pulseless Ventricular Tachycardia

#### Assessment

Ventricular Fibrillation, Ventricular Tachycardia  
Pulseless, apneic  
Confirm and record cardiac rhythm with electrodes verified in two leads on monitor

#### EMR and EMT

1. AED
2. CPR appropriate for patient's age
3. Oxygen 100% and airway maintenance appropriate to the patient's condition
4. **EMR and EMT STOP**

#### AEMT

1. IV NS TKO
2. **AEMT STOP**

#### PARAMEDIC

5. Epinephrine 1:10,000 1 mg IVP/IO q 4 mins (*peds Epinephrine 1:10,000 0.01 mg/kg IV/IO q 4 min*).
6. Defibrillate @ 200j biphasic, if no change in rhythm perform two minutes of CPR and evaluate rhythm. If no change in rhythm, repeat defibrillation, perform two minutes of CPR and evaluate rhythm. If no change in rhythm, continue 5 cycles of CPR then defibrillation cycle. (*peds begin at 2 j/kg*).
7. Administer:
  - a. Amiodarone 300 mg IV or IO, repeat after 5 min at 150 mg (*peds 5 mg/kg*)
  - b. For prolonged resuscitation or known acidosis consider: Sodium Bicarbonate 1 mEq/kg IV/IO followed by 0.5 mEq/kg q 10 min (*peds 1 mEq/kg may repeat at 0.5 mEq/kg q 10 min*)
8. CaCL 500 mg IVP (*peds 20 mg/kg*) if arrest secondary to renal failure, or history of hemodialysis.
9. Magnesium Sulfate 1-2 gm slow IV push over two minutes (*no pediatric dosage*).

#### NOTES:

- Defibrillation should not be delayed for any reason other than rescuer or bystander safety.

Prompt defibrillation is the major determinant of survival. Time on scene should be taken to aggressively treat ventricular fibrillation. Consider transport of patient after performing

# Roane County OES, EMS Division

## CARDIAC EMERGENCY

CPR/defibrillation cycles, securing the airway, obtaining IV/IO access, and administering two rounds of drugs. Remember, “stay and play” is the best chance for obtaining ROSC.

### 112 Ventricular Tachycardia with a Pulse

#### Assessment

Confirm and record cardiac rhythm with electrodes in two leads  
Check for palpable carotid pulse  
Decreased/altered mental status  
Dyspnea  
Chest pain/discomfort, suspected AMI  
Hypotension (systolic BP <90 mmHg)  
CHF/pulmonary edema  
Heart rate >150/min and QRS >.12 sec (3 small blocks)

#### EMR

1. Oxygen 100% and airway maintenance appropriate to the patient’s condition
2. Supportive care
3. **EMR STOP**

#### EMT

4. Pulse oximetry
5. **EMT STOP**

#### AEMT

6. Glucose check
7. INT or IV NS TKO
8. Titrate **Dextrose** 50% PRN slowly until normal levels achieved, if hypoglycemic.
9. **AEMT STOP**

#### PARAMEDIC

10. EKG monitor, 12 lead EKG
11. If rhythm is stable, regular and monomorphic admin **Adenosine** 12 mg Rapid IV Push
12. If systolic BP <90 mmHg **UNSTABLE**, prepare for synchronized cardioversion
  - A. Administer sedative as necessary, **Versed** 2-5 mg IV (*pediatrics 0.1 mg/kg*) and/or **Morphine** 2-5 mg.
  - B. Synchronize cardiovert beginning at manufacturers recommended initial energy setting, until heart rate <150/min (*pediatric begin at 0.5 j/kg*).
  - C. If rhythm converts, monitor for changes, transport. If rhythm does not convert, administer **Amiodarone** 150 mg over 10 minutes (*pediatrics 5 mg/kg*). Reattempt cardioversion.
  - D. Contact Medical Control

## Roane County OES, EMS Division

---

### CARDIAC EMERGENCY

13. If systolic BP >90 mmHg - **STABLE/ASYMPTOMATIC**
  - A. Have patient perform Valsalva maneuver for 10 seconds and administer **Amiodarone 150mg (pediatrics 5 mg/kg)** over 10 minutes.
  - B. If rhythm converts, monitor for changes, transport. If rhythm does not convert, administer **Amiodarone 150 mg** over 10 minutes (maximum three 150 mg doses) (pediatrics three doses of 5 mg/kg)

## CARDIAC EMERGENCY

### 113 Post Resuscitation

#### Assessment

Completion of arrhythmia treatment,  
Return of Spontaneous Circulation (ROSC)

1. Oxygen 100% and airway maintenance appropriate to the patient's condition

#### EMR and EMT

2. Supportive care
3. **EMR and EMT STOP**

#### AEMT

4. IV NS TKO
5. Assess BP - If systolic <90 mmHg administer 250 ml NS bolus (**pediatrics systolic BP 70 + 2x age in years, 20 ml/kg bolus**) repeat until BP >90 or appropriate for pediatric age.
6. Glucose check
7. Titrate **Dextrose 50%** prn slowly until normal levels achieved, if hypoglycemic.
8. **AEMT STOP**

#### PARAMEDIC

9. 12 Lead EKG
10. Medications:
  1. If anti-arrhythmic administered:
    - i. **Amiodarone** - 300 mg IV (**pediatrics 5 mg/kg, may repeat x3**), if one dose given and arrhythmia persist, give second dose of 150 mg
11. Continue ventilatory support to maintain ETCO<sub>2</sub> > 20, Respirations <12 ideally
12. Initiate induced hypothermia protocol if appropriate
13. If the patient does not tolerate the ET tube you may use **Versed** 2-5 mg IV (**pediatrics 0.1 mg/kg**) for patient sedation.

**Note:** Use of soft restraints if necessary for patient safety (to prevent extubation)

# Roane County OES, EMS Division

---

## ENVIRONMENTAL EMERGENCY

### 201 Chemical Exposure

**Special Note:** Personnel safety is the highest priority. **Do not handle the patient unless they have been decontaminated.** All EMS treatment should occur in the Support Zone after decontamination of the patient. Appropriate PPE will be utilized.

#### Assessment

History of exposure to chemical  
Identify substance and verify with documentation if possible  
Material Safety Data Sheets (M.S.D.S.) if available  
Stay within the appropriate zone for protection

**EMR EMT AEMT PARAMEDIC**

1. Oxygen and airway maintenance appropriate to patient's condition
2. Supportive care
3. IV NS TKO or INT PRN
4. Treatment - Standing Order If Internal Exposure and Conscious:
  1. Treat as Drug  
Ingestion
  2. Contact Medical  
Control **If External Exposure:**
    1. Remove victims clothing, jewelry, glasses, and contacts
    2. Decontaminate - EMS personnel must be wearing proper protective clothing prior to helping with the decontamination process.

#### Powder or like substance

1. Brush off patient
2. Flush with copious amounts of water for at least 20 minutes, assess for hypothermia q 5 min
3. Transport and continue flushing if necessary and if possible

#### Liquid substance

1. Flush with copious amounts of water for at least 20 minutes, assess for hypothermia q 5 min
2. Transport and continue if necessary and if possible **If**

#### **Inhalation:**

1. Reconsider Self-contained Breathing Apparatus
2. Remove victim from source ensuring there is no danger to personnel
3. Oxygen 100% and airway maintenance appropriate to patient's condition

#### **If Ocular:**

1. Immediately flush eye with tap water or normal saline for 15 minutes
2. Contact Medical Control

**NOTE:** Consult with HazMat scene IC prior to transport to ensure proper treatment and decontamination.

# Roane County OES, EMS Division

---

## ENVIRONMENTAL EMERGENCY

### 202 Drug Ingestion

#### Assessment

History of drug ingestion Level of consciousness (Alert, Verbal, Pain, or Unresponsive) Neurological status (LOC, pupils) General appearance (sweating, dry or flushed skin, signs of trauma)
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

#### EMR

1. Oxygen and airway maintenance appropriate to patient's condition
2. Ensure personnel protection from toxin and/or unruly patient
3. Supportive care
4. **EMR STOP**

#### EMT

5. Pulse oximetry
6. **EMT STOP**

#### AEMT

7. Glucose check
8. IV NS TKO or INT PRN
9. Titrate **Dextrose 50%** PRN slowly until normal levels achieved. (pediatrics 2 cc/kg D<sub>25</sub> IVP).
10. **Narcan 0.4 mg** IV/IO/IM/IN titrated to adequate ventilation (pediatrics 0.1 mg/kg IVP/IN) if narcotic use is suspected
11. **AEMT STOP**

#### PARAMEDIC

12. EKG monitor, 12 lead EKG
13. Consider **Versed 2-5 mg** IVP (pediatrics 0.1 mg/kg IVP) if patient is having seizures. May utilize **Versed IM** as per Seizure protocol based on weight if no IV present when seizure occurs.

**NOTES:** Poison Control may be contacted for **INFORMATION ONLY**. Treatment modalities are given within these protocols. Further treatments will be received through Medical Control.

## ENVIRONMENTAL EMERGENCY

### 203 Electrocution / Lightning Injuries

#### Assessment

Presence of signs and symptoms of electrical injury Entry / exit wounds

#### EMR

1. Oxygen and airway maintenance appropriate to the patient's condition
2. Spinal protection if electrocution/lightning over 1000 volts or suspicion of spinal injury
3. Supportive care
4. Treat burn per burn protocol
5. **EMR STOP**

#### EMT

6. Control any gross hemorrhage and dress wounds
7. Pulse oximetry
8. **EMT STOP**

#### AEMT

9. IV LR if signs of shock 20 cc/kg bolus of fluid (peds 20 cc/kg bolus)
10. **AEMT STOP**

#### PARAMEDIC

11. EKG Monitor, 12 Lead EKG
12. Consider 2<sup>nd</sup> IV en route to hospital
13. Consider pain medications
  - A. Morphine 2-10 mg/IV/IO (peds 0.1 mg/kg IV/IO)
  - B. Fentanyl 25-50 mcg/IV/IO (peds 0.1 mcg/kg IV/IO)

# Roane County OES, EMS Division

## ENVIRONMENTAL EMERGENCY

### 204 Hyperthermia

#### Assessment

History of exposure to warm temperature / Febrile  
May have hot and dry or warm and moist skin  
May be hypotensive  
Determine history of therapeutic drug use; history of substance abuse  
Poor skin turgor  
Signs of hypovolemic shock  
History of infection or illness  
Dark urine - Suggest muscle breakdown and possible kidney damage  
Tachycardia, Hyperventilation, Hypertension  
Neurologic - Light headedness, confusion to coma, seizures

#### EMR

1. Oxygen and airway maintenance appropriate to the patient's condition
2. Remove clothing, apply wet linen or wet abdominal pads to groin and axillary areas
  - A. Expose to circulating air
  - B. **DO NOT** cool patient to the point of shivering
3. Move patient to protected environment (shade, AC, etc.)
4. **EMR STOP**

#### EMT

5. Pulse oximetry
6. **EMT STOP**

#### AEMT

7. Glucose check
8. IV NS or LR 20 cc/kg bolus (*peds 20 cc/kg bolus*)
  - A. Repeat second bolus of fluids if needed
  - B. Oral rehydration if patient able to maintain airway
  - C. IV NS - rate proper for patient condition
9. GENTLY massage extremities to prevent cold induced vasoconstriction
10. Titrate Dextrose 50% PRN slowly until normal levels achieved, if hypoglycemic
11. **AEMT STOP**

#### PARAMEDIC

12. EKG Monitor, 12 Lead EKG if cardiac effects considered
  - A. Time is of the essence in decreasing the patient's body temperature
  - B. Do not use IV iced saline for cooling patient.
  - C. Hyperthermia may be cause by one of the following:
    - I. Antipsychotic medications and major tranquilizers: Phenothiazine (Thorazine®), Butyrophenones (Haldol®)

## Roane County OES, EMS Division

---

### ENVIRONMENTAL EMERGENCY

- II. Cyclic antidepressants such as: Elavil®, Norpramin®, Tofranil®
- III. Amphetamines
- IV. Monoamine oxidase inhibitors such as: Nardil®, Marplan®
- V. Anticholinergic drugs such as: Atropine, Congentin, Scopolamine
- VI. Illicit drugs: Cocaine, PCP, LSD, Ecstasy (MDMA)

# Roane County OES, EMS Division

## ENVIRONMENTAL EMERGENCY

### 205 Hypothermia

#### Assessment

History of exposure to cold temperature including duration  
Core body temperatures  $<92^{\circ}$   
Drug/Alcohol use  
CNS Depressants  
Examine for associated trauma  
Immersion in cold water  
Predisposing medical condition  
Signs: Vital signs, Bradycardia, Hypotension, Cold extremities, Neurologic (confusion, altered LOC, coma)

#### EMR and EMT

1. Oxygen 100% at 12-15 Lpm with BVM
2. Remove the patient from the cold environment
3. Remove wet clothing and cover with warm, dry blankets
4. Evaluate pulse for one full minute (Do not perform CPR until NO PULSE is confirmed)
5. Handle patient gently (*aggressive movement may trigger V-Fib*)
6. Do not allow patient to walk or exert themselves
7. Do not massage extremities
8. **EMR and EMT STOP**

#### AEMT

9. Glucose check
10. IV NS 75 cc/hr warmed if possible (*peds 4 cc/kg/hr max 150 cc/hr*)
11. Titrate Dextrose 50% PRN slowly until normal levels achieved, if hypotensive
12. If patient in coma, Narcan 0.4 mg IV/IO/IM/IN titrated to adequate ventilation (*peds 0.1 mg/kg slow IVP/IN*)
13. **AEMT STOP**

#### PARAMEDIC

14. EKG Monitor, No CPR if bradycardic rhythm exists
  15. If body temperature  $>85^{\circ}$  F - follow normal arrest protocols 16.  
If body temperature  $<85^{\circ}$  F and patient in V-Fib:
    - A. Defibrillate @ 100j, begin CPR defibrillate at 2 min intervals, increase joules at each interval until 200j max (120j, 150j, 200j) (*peds 2 j/kg then 4 j/kg*)
    - B. Withhold meds and further shocks until patient warmed to  $>85^{\circ}$  F
    - C. Continue CPR and rewarming attempts **NOTES:**
      - If patient is alert and responding appropriately, rewarm actively:
      - Heat packs or warm water bottles to the groin axillary and cervical areas
      - If patient is unresponsive, rewarm passively:
      - Increase the room temperature gradually, cover with blankets
- CONTINUED ON NEXT PAGE

### ENVIRONMENTAL EMERGENCY

- If the following are signs and symptoms found at varying body temperature:
  - 95° - amnesia, poor judgment, hyperventilation, bradycardia, shivering
  - 90° - loss of coordination, decreased resp., shivering ceases, bradycardia
  - 85° - decreased LOC, slow respirations, atrial fibrillation, decreased BP, decreased heart rate, ventricular irritability

## ENVIRONMENTAL EMERGENCY

### 206 Near Drowning

#### Assessment

History compatible with near drowning  
Suspect hypothermia in “cold water” near drowning Suspect cervical spine injury

#### EMR and EMT

1. Oxygen and airway maintenance appropriate to patient’s condition  
The Heimlich Maneuver may be indicated for airway obstruction  
Gastric decompression may be necessary to ensure adequate respirations or ventilations  
If necessary, ventilations may be started prior to patient’s removal from water
2. Remove patient from water, clear airway while protecting the C-spine ASAP
3. ***If patient is unconscious and pulseless - refer to the Cardiac Arrest Protocol***
4. ***If Hypothermic - go to hypothermia protocol***
5. Supportive care
6. **EMR and EMT STOP**

#### AEMT

7. INT or IV NS TKO, if hypotensive give  
20 cc/kg bolus of fluid (***peds 20 cc/kg***)
8. **AEMT STOP**

#### PARAMEDIC

9. EKG Monitor and treatment specific for the arrhythmia

#### NOTES:

- Reinforce the need to transport and evaluation for all patients with a submersion incident.
- Consider C-Spine protection.

# Roane County OES, EMS Division

## ENVIRONMENTAL EMERGENCY

### 207 Nerve Agent Exposure

**Special Note:** Personnel safety is the highest priority. Do not handle the patient unless they have been decontaminated. All EMS treatment should occur in the Support Zone after decontamination of the patient. Appropriate PPE will be utilized.

#### Assessment

History of exposure  
Hyper-stimulation of muscarinic sites (smooth muscles, glands) and nicotinic sites (skeletal muscles, ganglions)  
Increased secretions: Saliva, tears, runny nose, secretions in airways, secretions in GI tract, sweating  
Pinpoint pupils  
Narrowing airway  
Nausea, vomiting, diarrhea  
Fasciculations, Flaccid paralysis, general weakness  
Tachycardia, hypertension  
Loss of consciousness, convulsions, apnea

#### EMR

1. Oxygen and airway maintenance appropriate to the patient's condition
2. Depending on signs and symptoms administer Nerve Agent Antidote kit
  - A. Mild - Increased secretions, pinpoint pupils, general weakness
    - I. Decontamination, supportive care
  - B. Moderate/Severe - mild symptoms, dyspnea, unconscious, convulsions, apnea
    - II. 1 Nerve Agent antidote kit
    - III. May be repeated in 5 min, prn
3. Keep patient warm
4. **EMR STOP**

#### EMT

5. Pulse oximetry
6. **EMT STOP**

#### AEMT

7. IV NS TKO
8. **AEMT STOP**

#### PARAMEDIC

9. EKG Monitor, 12 Lead EKG if cardiac effects considered
10. **Versed 2-5 mg IV for seizures (peds 0.1 mg/kg)**

#### Treatment - Protocol:

Repeated doses of Atropine may be required after Nerve Agent Antidote Kit(s) given  
**CONTINUED ON NEXT PAGE**

# Roane County OES, EMS Division

---

## ENVIRONMENTAL EMERGENCY

### NOTES:

- This is for mass casualty situations and is dependent on supplies available. There is no contraindication for the use of a Nerve Agent Antidote Kit in the case of true nerve agent exposure.

# Roane County OES, EMS Division

## ENVIRONMENTAL EMERGENCY

### 208 Poisonous Snake Bite

#### Assessment

Protect yourself from the exposure of snakebite. Snakes can envenomate up to one hour after death.  
Determine type of snake if possible, time of bite, and changes in signs and symptoms since occurrence.  
If possible, transport the **DEAD** snake in a secured vessel with the victim for identification  
Paresthesias (numbing or tingling of mouth, tongue, or other areas)  
Local pain  
Peculiar or metallic taste  
Chills, nausea and vomiting, headache, dysphagia  
Hypotension  
Fever  
Local edema, blebs (blister or pustule jewel), discoloration Bite wound configuration

#### EMR

1. Remove rings and bracelets from the patient
2. Oxygen and airway maintenance appropriate to patient's condition
3. Immobilize affected area keeping extremities in neutral position
4. Mark progression of swelling at the time of initial assessment and q 5 minutes
5. Supportive care
6. **EMR STOP**

#### EMT

7. Pulse oximetry
8. **EMT STOP**

#### AEMT

9. INT or IV NS TKO, if hypotensive 20 cc/kg (*peds 20 cc/kg*)
10. **AEMT STOP**

#### PARAMEDIC

11. EKG Monitor, 12 Lead EKG if cardiac effects considered

#### Treatment - Protocol:

**Versed** may be indicated if anxiety is overwhelming. Contact Medical Control prior to initiating therapy.

#### NOTE:

- DO NOT USE ice, tourniquets, hemorrhage control clamp or constricting bands at the bite site or proximal to bite site. If already applied, remove.
- Do NOT place IV in affected extremity if possible.

# Roane County OES, EMS Division

---

## ENVIRONMENTAL EMERGENCY

### 209 Radiation/Hazmat

**Special Note:** Personnel safety is the highest priority. **Do not handle the patient unless they have been decontaminated.** All EMS treatment should occur in the Support Zone after decontamination of the patient. Appropriate PPE will be utilized.

#### Assessment

Extent of radiation/chemical exposure (number of victims, skin vs. inhalation exposure) Nature of exposure Symptoms exhibited by patient Neurologic status (LOC, pupil size) General appearance (dry or sweaty skin, flushed, cyanotic, singed hair) Associated injuries Decontamination prior to treatment
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

#### EMR

1. If eye exposure, irrigate for a minimum of 20 minutes
2. Treat associated injuries (LSB, limb immobilization, wound treatment)
3. Supportive care
4. Treat per burn protocol
5. Oxygen and airway maintenance appropriate to the patient's condition
6. **EMR STOP**

#### EMT

7. Pulse oximetry (keep sats >94%)
8. **EMT STOP**

#### AEMT

9. INT or IV NS/LR, if hypotensive 20 cc/kg (*peds 20 cc/kg*)
10. **AEMT STOP**

#### PARAMEDIC

11. EKG Monitor, 12 Lead EKG if cardiac cause considered

**NOTE:** Consult with HazMat scene IC prior to transport to ensure proper treatment and decontamination.

# Roane County OES, EMS Division

---

## ENVIRONMENTAL EMERGENCY

### 210 Carbon Monoxide Exposure

#### Assessment

Known or suspected CO exposure (Active fire scene) Suspected source/duration exposure Known or possible pregnancy Measured atmospheric levels Past medical history, medications Altered mental status/dizziness Headache, Nausea/vomiting Chest pain/respiratory distress Neurological impairments Vision problems/reddened eyes Tachycardia/tachypnea Arrhythmias, seizures, coma
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

#### **EMR EMT AEMT PARAMEDIC**

#### Measure Carbon Monoxide COHb % (SpCO)

- A. If SpCO is 0%-5% nor further medical evaluation of SpCO is required\*
- B. SpCO <15% and SpO<sub>2</sub> >90%
  - I. If patient has NO symptoms of CO and/or Hypoxia treatment for CO exposure is required\*
  - II. Recommend evaluation of home/work environment for presence of CO
- C. SpCO >15%
  - I. Oxygen via NRB and transport to ED
  - II. If cardiac/respiratory/neurological symptoms go to the appropriate protocol

#### NOTES:

- If monitoring responders at fire scene, proceed with Scene Rehabilitation Protocol where applicable.
- \*Fetal hemoglobin has a greater attraction for CO than maternal hemoglobin. Females who are known to be pregnant or who could be pregnant should be advised that EMS measured SpCO levels reflect the adult's level, and that fetal COHb levels may be higher. Recommend transport for a hospital evaluation for any CO exposed pregnant person.
- The absence (or low detected levels of COHb is not a reliable predictor of firefighter or victim exposure to other toxic byproducts of fire.
- In obtunded fire victims, consider HazMat Cyanide treatment protocol.

The differential list for CO toxicity is extensive. Attempt to evaluate other correctable causes when possible.

## MEDICAL EMERGENCIES

### 300 Medical Complaint Not Specified under other Protocols

#### Assessment

Pertinent history to complaint  
Allergies/Medications taken or prescribed  
Provocation  
Quality of Pain / Discomfort  
Relieved by  
Signs and symptoms  
Onset, type, and duration of pain

#### EMR

1. Oxygen and airway maintenance appropriate for the patient's condition
2. Patient positioning appropriate for condition
3. Supportive care
4. **EMR STOP**

#### EMT

5. Pulse oximetry
6. **EMT STOP**

#### AEMT

7. Glucose check
8. Titrate **Dextrose** 50% PRN slowly until normal levels achieved. If indicated, INT or IV NS TKO unless signs of shock, then 20 cc/kg fluid bolus
9. **AEMT STOP**

#### PARAMEDIC

10. EKG Monitor, 12 Lead EKG if cardiac cause considered

## 301 Abdominal Pain (non-traumatic)/Nausea and Vomiting

### Assessment

Description of pain, onset, duration, location, character, radiation  
Aggravating factors, last menstrual periods in females, vaginal bleeding in females  
Recent trauma  
History of abdominal surgery or problems  
Blood in urine, vomitus, or stool  
Nausea, vomiting, diarrhea  
Fever, diaphoresis, jaundice  
Abdomen: tenderness, masses, rigidity, hernia, pregnancy, distension, guarding

### EMR

1. Oxygen and airway maintenance appropriate to the patient's condition
2. Allow patient to assume comfortable position or place patient supine, with legs elevated with flexion at hip and knees unless respiratory compromise or a procedure contraindicates
3. Supportive care
4. **EMR STOP**

### EMT

5. Pulse oximetry
6. **EMT STOP**

### AEMT

7. IV NS 20 cc/kg, if signs of shock (peds 20 cc/kg bolus)
8. **AEMT STOP**

### PARAMEDIC

9. EKG Monitor, 12 Lead EKG if considering cardiac causes
10. Ondansetron (Zofran) 2-4mg IV (peds 0.15mg/kg IV) if intractable nausea and persistent vomiting and no signs of shock. Use lower dose initially especially in the elderly.
11. If allergic to Ondansetron, Phenergan 6.25mg -12.5mg IVP or IM, only if no signs or symptoms of shock. Use lower dose initially, especially in the elderly. May repeat once if needed.
12. Consider second IV en route if patient exhibits signs of shock

## 302 Acute Pulmonary Edema / CHF

### Assessment

Focus assessment of Airway, Breathing, and Circulation  
Shortness of breath  
Cyanosis  
Pedal Edema  
Profuse sweating, or cool and clammy skin  
Erect posture  
Distended neck veins (engorged, pulsating) - late sign  
Bilateral rales/wheezes  
Tachycardia (rapid pulse >100 bpm)  
History of CHF or other heart disease, or renal dialysis Lasix or Digoxin on medication list

### EMR

1. Oxygen and airway maintenance appropriate to patient's condition. If respiration is <10/min, or >30/min, consider assisted ventilations w/ BVM and 100% Oxygen
2. Keep patient in upright seated position
3. **EMR STOP**

### EMT

4. If the patient has Albuterol Inhalation Treatment prescribed, assist them with one treatment.
5. If Systolic BP is >110 and the patient is symptomatic, **assist** patient with 1 NTG dose sublingually and reassess every 5 minutes. (Refer to the medication assist procedure) Maximum of three doses. Use caution in patients taking erectile dysfunction medications. Profound hypotension may occur.
6. **EMT STOP**

### AEMT

7. INT
8. If Systolic BP >90 mmHg
  - a. Assess for crackles, wheezes, or rales, JVD, peripheral edema, cyanosis, diaphoresis, respiratory rate >25/min or <10/min then:
    - i. One NTG tablet SL. Repeat NTG q 5 min after initial dose. **STOP** therapy if systolic BP <100 mmHG;
    - ii. **Albuterol** 2.5mg/3 ml via nebulizer q 5 minutes, to max of 3 doses;
9. If Systolic BP <90 mmHg
  - b. Continue oxygen and initiate rapid transport, see hypotension protocol, contact  
Medical Control immediately
10. **AEMT STOP**  
(CONTINUED ON NEXT PAGE →)

PARAMEDIC

11. EKG Monitor, 12 lead performed and transmitted if needed
12. If severe respiratory distress and no contraindications. Begin CPAP
13. If patient is hypotensive, **Dopamine** 400 mg/250 cc D5W IV mix, begin at 15 cc/hr titrate up, systolic pressure <90 mmHg and symptomatic is considered hypotensive.

## 303 Anaphylactic Shock

### Assessment

Contact with a known allergen or with substances that have a high potential for allergic reactions  
Sudden onset with rapid progression of symptoms  
Dyspnea, presents with an audible wheeze on confrontation, generalized wheeze on auscultation, decreased air exchange on auscultation  
Generalized urticaria, erythema, and angioedema especially noticeable to face and neck  
Complaint of chest tightness or inability to take a deep breath

### EMR

1. Position of comfort, reassure
2. Oxygen and airway maintenance appropriate for patient's condition, pulse oximetry
3. **EMR STOP**

### EMT

4. Pulse oximetry
5. If patient has a prescribed Epinephrine for Anaphylaxis, assist patient with administration
6. **EMT STOP**

### AEMT

7. Epinephrine 1:1,000 0.3 mg IM or 1:10,000 IV/IO (pediatrics Epinephrine 1:1,000 0.01 mg/kg IM or 1:10,000 IV/IO, max dose is 0.3 mg)
8. Albuterol 2.5 mg/3ml, nebulizer treatment if wheezing is present
9. IV NS or LR, large bore @ TKO - If hypotensive 20 cc/kg bolus (pediatrics 20 cc/kg bolus)
10. **AEMT STOP**

### PARAMEDIC

11. EKG Monitor, 12 Lead EKG if cardiac cause considered
12. Diphenhydramine (Benadryl) 25-50 mg IV/IO or deep IM (pediatrics 1 mg/kg IVP)
13. Solu-Medrol 125 mg IVP (pediatrics contact Medical Control)

## 304 Cerebrovascular Accident (CVA)

### Assessment

Altered Level of consciousness (coma, stupor, confusion, seizures, delirium)  
Intense or unusually severe headache of sudden onset or any headache associated with decreased level of consciousness or neurological deficit, unusual / severe neck or facial pain  
Aphasia/Dysphasia (unable to speak, incoherent speech, or difficulty speaking)  
Facial weakness or asymmetry (paralysis of facial muscles, usually noted with the patient speaks or smiles); may be on the same side or opposite side from limb paralysis  
In-coordination, weakness, paralysis, or sensory loss of one or more limbs; usually involves one half of the body particularly the hand  
Ataxia (poor balance, clumsiness, or difficulty walking)  
Visual loss (monocular or binocular); may be a partial loss of visual field  
Intense vertigo, double vision, unilateral hearing loss, nausea, vomiting, photophobia, or phonophobia

### EMR

1. Oxygen and airway maintenance appropriate for the patient's condition
2. Continually monitor airway due to decreased gag reflex and increased secretions
3. Conduct a brief targeted history and physical exam. Establish time of onset. Document witness to Time of Onset and their contact information. Include the Cincinnati Pre-Hospital Stroke Scale.
4. **EMR STOP**

### EMT

5. Maintain body heat, protect affected limbs from injury, anticipate seizures
6. If seizure present follow seizure protocol
7. If shock signs present follow shock protocol
8. If trauma suspected, spinal stabilization. Elevate head 30° if no of spinal injury
9. Pulse oximetry
10. **EMT STOP**

### AEMT

11. Glucose check
12. IV NS TKO (30 cc/hr) or INT
13. Titrate **Dextrose** 50% PRN slowly until normal levels achieved, if hypoglycemic
14. **Narcan** 0.4 mg IV/IO/IM/IN titrated to adequate ventilation (*peds 0.1 mg/kg slow IVP/IN*) May repeat dose up to 2 mg if narcotics suspected
15. **AEMT STOP**

CONTINUED ON NEXT PAGE

## PARAMEDIC

16. EKG Monitor, 12 Lead EKG
17. Complete thrombolytic screening protocol
18. Complete Stroke assessment scale
19. If positive for CVA, recommend transport to stroke center
20. Contact Medical Control if SBP >220 or DBP>140 if authorized to give Nitro spray q 5 min. Goal is to reduce blood pressure by 15%

### REFERENCE-CVA

#### THE CINCINNATI PREHOSPITAL STROKE SCALE

**Facial Droop** (have patient show teeth or smile):

Normal: Both sides of face will move equally well.

Abnormal: One side of face does not move as well as the other side

**Arm Drift** (patient closes eyes and holds both arms out):

Normal: Both arms move the same or both arms do not move at all.

Findings, such as pronator grip may be helpful

Abnormal: One arm does not move or one arm drifts down compared with the other

**Speech** (have the patient say “you can’t teach an old dog new tricks):

Normal: Patient uses correct words with no slurring

Abnormal: Patient slurs word(s), uses inappropriate words, or is unable to speak

For evaluation of acute, non-comatose, non-traumatic neurologic complaint.

<b>Facial/Smile or Grimace:</b>		
Have the patient show teeth or smile.	Normal: Both sides of the face move equally	Abnormal: Left or right side of face does not move as well
<b>Arm Drift:</b>		
Have the patient close both eyes and hold both arms straight out for 10 seconds	Normal: Arms move equally or do not move	Abnormal: Left or right arm does not move or drifts down
<b>Speech:</b>		
Have the patient repeat a simple phrase such as “It is sunny outside today”	Normal: Words stated correctly without slurring	Abnormal: Patient slurs words or uses the wrong words, or is unable to speak.

## Roane County OES, EMS Division

---

### PREHOSPITAL SCREEN FOR THROMBOLYTIC THERAPY

Complete this report for all patients symptomatic for Acute Coronary Syndrome or CVA. Report to the Emergency Department Physician/Nurse any positive findings. Document all findings in the PCR.

Witness/next of kin contact info: _____ _____
-----------------------------------------------------

Time of onset of the symptoms: _____ _____
--------------------------------------------------

Systolic BP >240 mmHg	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Diastolic BP >110 mmHg	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Right arm vs. Left arm Systolic BP difference >15 mmHg	<input type="checkbox"/> Yes	<input type="checkbox"/> No
History of recent brain/spinal cord surgery, CVA, or injury	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Recent trauma or surgery	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Bleeding disorder that causes the patient to bleed excessive	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Prolonged CPR (>10 minutes)	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Pregnancy	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Taking Coumadin, Aspirin, or other blood thinners	<input type="checkbox"/> Yes	<input type="checkbox"/> No

## 305 Croup

### Assessment

History - Viral infections resulting in inflammation of the larynx, trachea  
Seasonal - Late fall/early winter  
Children under 6 yrs old with cold symptoms for 1-3 days  
Hoarseness  
Barking, Seal-like cough  
Stridor, NOT wheezes  
Low grade fever  
No history of obstruction, foreign body, trauma

### EMR and EMT

1. Oxygen and airway maintenance appropriate to the patient's condition
2. Allow patient to assume comfortable position or place patient supine
3. Supportive care
4. **EMR and EMT STOP**

### AEMT and PARAMEDIC

5. Nebulized Epinephrine 1:1000
  - A. 1 mg diluted to 2.5-3 cc with saline flush, nebulized (mask or blow-by)
  - B. May repeat up to 3 total doses
  - C. If the patient has significant distress, 3 ml (3 mg) diluted with 2.5 to 3 cc saline flush may be administered as initial aerosol
6. Contact Medical Control for subsequent aerosols.

# Roane County OES, EMS Division

---

## MEDICAL EMERGENCY

### 306 Family Violence

#### Assessment

Fear of household member Reluctance to respond when questioned Unusual isolation, unhealthy, unsafe living environment Poor personal hygiene/inappropriate clothing Conflicting accounts of the incident History inconsistent with injury or illness Indifferent or angry household member Household member refused to permit transport Household member prevents patient from interacting openly or privately Concern about minor issues but not major ones Household with previous violence Unexplained delay in seeking treatment
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

#### **EMR EMT AEMT PARAMEDIC**

\*Direct questions to ask when alone with patient and time available:

1. Has anyone at home ever hurt you?
2. Has anyone at home touched you without your consent?
3. Has anyone ever made you do things you didn't want to do?
4. Has anyone taken things that were yours without asking?
5. Has anyone scolded or threatened you?
6. Are you afraid of anyone at home?

\*\*Signs and Symptoms

- Injury to soft tissue areas that are normally protected
- Bruise or burn in the shape of an object
- Bite marks
- Rib fracture in the absence of major trauma
- Multiple bruising in various stages of healing

#### Treatment - Standing Order

1. Patient care is first priority
2. If possible remove patient from situation and transport
3. Police assistance as needed
4. If sexual assault follow sexual assault protocol
5. Obtain information from patient and caregiver
6. Do not judge
7. Report suspected abuse to hospital after arrival. Make verbal and written report.

**NOTE:** National Domestic Violence Hotline 1 (800) 799- SAFE (7233)

## MEDICAL EMERGENCY

### 307 Hyperglycemia Associated with Diabetes

#### Assessment

History of onset  
Altered level of consciousness  
Pulse: tachycardia, thready pulse  
Respirations (Kussmaul-Kien - air hunger)  
Hypotension  
Dry mucous membranes  
Skin may be cool (consider Hypothermia)  
Ketone odor on breath (Acetone smell)  
Abdominal pain, nausea and vomiting  
History of polyuria or polydipsia (excessive urination or thirst) Blood glucose determination

#### EMR

1. Oxygen and airway maintenance appropriate to patient's condition
2. Supportive care
3. **EMR STOP**

#### EMT

4. Suction airway as needed.
5. Pulse oximetry
6. **EMT STOP**

#### AEMT

7. Glucose check
8. IV NS TKO or INT. Consider 250-500 cc NS bolus, only in patients with signs of dehydration, vomiting or DKA
9. If BS >250 mg/dL, start 10-20 cc/kg infusions of NS (*peds 4 cc/kg/hr max 150 cc/hr DO NOT bolus*), then reassess blood sugar
10. **AEMT STOP**

#### PARAMEDIC

11. EKG Monitor, 12 Lead EKG if cardiac cause considered

# Roane County OES, EMS Division

## MEDICAL EMERGENCY

### 309 Hypoglycemia

#### Assessment

History of onset of event  
History of Insulin excess (overdose, missed meal, exercise, vomiting, or diarrhea)  
Confusion, agitation, headaches, or comatose  
Pulse rate (normal to tachycardia)  
Respirations (shallow, slow)  
Skin (sweaty, often cool)  
Flaccid muscle tone  
Grand Mal seizure  
Fecal, urinary incontinence

#### EMR

1. Oxygen and airway maintenance appropriate to patient's condition (snoring respirations is a sign of an INADEQUATE airway)
2. Supportive care
3. **EMR STOP**

#### EMT

4. If patient is known diabetic and is conscious with an intact gag reflex, administer one tube of instant Glucose and reassess
5. Pulse oximetry
6. **EMT STOP**

#### AEMT

7. Glucose check
8. IV NS TKO
9. If blood sugar is less than 80 mg/dL and symptomatic: administer 25 grams of Dextrose, (*peds 2 cc/kg D<sub>25</sub> IV/IO; if needed a mixture of D<sub>50</sub> and Normal Saline can be obtained through mixing 1 cc to 1 cc for the treatment of symptomatic hypoglycemia in pediatric patients*) reassess blood sugar level q 15 min, administer 10-25 grams Dextrose PRN
10. **AEMT STOP**

#### PARAMEDIC

11. EKG Monitor, 12 Lead EKG if cardiac cause considered

# Roane County OES, EMS Division

---

## MEDICAL EMERGENCY

### 310 Medications at Schools

To provide authorization for the use of medications not commonly used. **For emergency use only.**

Assessment

The patient must exhibit the signs and symptoms for which the medication is prescribed

#### EMR EMT AEMT PARAMEDIC

1. Oxygen and airway maintenance appropriate to the patient's condition
2. Other treatments will be in accordance with the EMS BLS/ALS SOPs
3. Necessary medication(s) administration as requested by school official(s)
  - A. Schools must provide the medication(s) to be administered
  - B. Schools must provide a written copy of the physician order and care plan for attachment to the patient care report
  - C. This documentation by the patient's primary physician should list the following:
    - I. Name of the patient
    - II. Name of the primary physician
    - III. Document must be signed by the primary physician
    - IV. Contact phone number of the primary physician
    - V. Name of medication(s)
    - VI. Signs and symptoms for which the medication(s) is/are prescribed
    - VII. Dosage of the medication(s)
    - VIII. Number of repeat doses of the medication(s)
    - IX. Route(s) of administration(s)
    - X. Potential side effects of the medication(s)
4. Medication(s) will only be administered if the patient meets the signs and symptoms for that medication
5. Copies of the care plan and physician order must be attached to the patient care report
6. If the medication(s) is/are not administered documentation must include reasons for withholding
7. Whenever medication is administered under these circumstances transport is mandatory

**NOTE:** If you have any additional questions or concerns please contact Medical Control.

# Roane County OES, EMS Division

---

## MEDICAL EMERGENCY

### 311 Non-Formulary Medications

To provide authorization for the use of medications not commonly used within the current guidelines. **For Emergency Use Only.**

Assessment

The patient must exhibit the signs and symptoms for which the medication is prescribed

#### EMR EMT AEMT PARAMEDIC

1. Oxygen and airway maintenance appropriate for patient's condition
2. Other treatment will be in accordance with the BLS/ALS SOPs
3. Necessary medication(s) administration as requested by caregiver(s):
  - A. Caregiver must provide the medication(s) to be administered
  - B. Caregiver must provide a written copy of the physician order and care plan for attachment to the patient care report
  - C. This documentation by the patient's physician should list the following:
    - I. Name of the patient
    - II. Name of primary physician
    - III. Document must be signed by the primary physician
    - IV. Contact phone number of the primary physician
    - V. Name of the medication(s)
    - VI. Signs and symptoms for which the medication(s) is prescribed
    - VII. Dosage of the medication(s)
    - VIII. Number of repeat doses of the medication(s)
    - IX. Route(s) of administration(s)
    - X. Potential side-effects of medication(s)
4. Medication(s) will only be administered if the patient meets the signs and symptoms for that medication.
5. Copies of the care plan and physician order must be attached to the patient care report
6. If the medication(s) is/are not administered documentation must include those reasons for withholding
7. Whenever medication is administered under these circumstances, transport is mandatory

**NOTE:** If you have any additional questions or concerns please contact Medical Control.

# Roane County OES, EMS Division

## MEDICAL EMERGENCY

### 312 Respiratory Distress (Asthma/COPD)

#### Assessment

**Mild attack** - Slight increase in respiratory rate. Mild wheezes. Good skin color. **Moderate attack** - Marked increase in respiratory rate. Wheezes easily heard. Accessory muscle breathing  
**Severe attack** - Respiratory rate more than twice normal. Loud wheezes or so tight no wheezes are heard, patient anxious. Grey or ashen skin color. Hx - COPD, Emphysema, Asthma, or other restrictive lung disease  
Respiratory rate greater than 25 per minute or less than 10 per minute  
Labored respiration, use of accessory muscles or tripodding  
Breath sounds: Bilaterally diminished, dry crackles, wheezing  
Cyanosis/Diaphoresis  
Use of short sentences  
Unilateral breath sounds

#### EMR

1. Oxygen and airway maintenance appropriate for patient's condition
2. **EMR STOP**

#### EMT

3. If the patient has prescribed **Albuterol** Inhalation treatment, assist the patient with 2.5 mg/ 3 ml NS and start the oxygen flow rate at 6 LPM or until the appropriate mist is achieved.
4. If patient uses a MDI, assist patient with one dose
5. **EMT STOP**

#### AEMT

6. Pulse oximetry and capnography
7. INT or IV NS TKO
8. Administer **Albuterol** 2.5 mg/3 ml NS and start the oxygen flow rate at 6 LPM or until the appropriate mist is achieved
9. **Epinephrine 1:1000 IM or 1:10,000 IV/IO 0.3-0.5 mg (*pediatrics 1:1000 0.01 mg/kg IM or 1:10,000 IV/IO, max dose is 0.3 mg*)** for patients in severe distress. Be mindful of cardiac side effects.
10. **AEMT STOP**

(CONTINUED ON NEXT PAGE)

## Roane County OES, EMS Division

---

### MEDICAL EMERGENCY

#### PARAMEDIC

11. EKG Monitor, 12 lead if cardiac cause considered
12. 1 **Atrovent** unit dose, 0.5 mg/2.5 ml, should be added to a single dose of **Albuterol** for nebulization if not already done by other Healthcare provider.
13. If no change in condition, consider continuous nebulizer of Albuterol up to 10 mg (4 total doses of **Albuterol** and 1 total dose of **Atrovent**)
14. In severe cases consider **Solu-Medrol** 125 mg IV (*pediatrics contact Medical Control*)
15. Use CPAP if no contraindications in conjunction with **Albuterol** nebulizer treatments

**\*\*\*Pediatrics: consult Medical Control prior to administering Solu-Medrol\*\*\***

# Roane County OES, EMS Division

## MEDICAL EMERGENCY

### 313 Seizures

#### Assessment

Seizure (onset, duration, type, post-seizure, level of consciousness)  
Medical (diabetes, headaches, drugs, alcohol, seizure history)  
Physical (seizure activity, level of consciousness, incontinence, head and mouth trauma, vital signs)  
Trauma (head injury or hypoxia secondary to trauma)

#### EMR

1. Oxygen and airway maintenance appropriate to patient's condition
2. Protect patient from injury during active seizures
3. C spine precautions if appropriate
4. **EMR STOP**

#### EMT

5. If febrile cool as per hyperthermia protocol and monitor
6. **EMT STOP**

#### AEMT

7. Glucose check
8. IV/IO NS TKO or INT
9. Titrate **Dextrose** 50% PRN slowly until normal levels achieved, if blood sugar <80 mg/dl
10. If narcotic overdose, **Narcan** 0.4 mg IV/IO/IM/IN titrated to adequate ventilation. (***pediatrics 0.1 mg/kg, titrated to adequate ventilation***). May repeat dose up to 2mg.
11. **AEMT STOP**

#### PARAMEDIC

12. EKG monitor- treat dysrhythmia per protocol
13. Adults - If actively seizing:
  - A. Before an IV is established, administer **Versed** 10 mg IM (>39 kg) or 5 mg IM (<39 kg) for active seizure.
  - B. If IV established, **Versed** 2-5 mg IV/IO may repeat if seizure continues
14. Pediatrics - If actively seizing:
  - A. **Versed 0.1 mg/kg IV/IO or if no IV Versed 0.15 mg/kg IM up to 5 mg**
  - B. **If seizure persists for 4 minutes repeat medication once**

CONTINUED ON NEXT PAGE

## Roane County OES, EMS Division

---

### MEDICAL EMERGENCY

#### NOTES:

- Specifically evaluate for: active bleeding, trauma, eye deviation, pupil equality, mouth or tongue bleeding, Urinary or fecal incontinence, lack of arm or leg movement or tone.
- The goal of Narcan therapy is to restore adequate ventilation. Larger doses especially in patients on chronic opiate therapy, often need very small doses in the event of overdose. Larger doses of Narcan usually create more agitation

# Roane County OES, EMS Division

---

## MEDICAL EMERGENCY

### 314 Sexual Assault

#### Assessment

Traumatic Injuries

#### **EMR EMT AEMT PARAMEDIC**

1. Oxygen and airway maintenance appropriate to patient's condition
2. Be calm and assuring with sensitivity toward the patient
3. DO NOT make unnecessary physical contact with the patient
4. If possible, have a witness the same gender as the victim present at all times
5. Wrap a plastic sheet around the victim if possible
6. DO NOT inspect genitals unless evidence of uncontrolled hemorrhage, trauma, or severe pain is present
7. DO NOT allow patient to shower or douche
8. Collect patient's clothing when possible
  - A. Place clothing in plastic sheet or separate plastic/paper bags with ID labels and found location
  - B. Leave all sheets placed in plastic/paper bag with patient at facility
  - C. Notify all staff of clothing samples
9. Transport patient to appropriate facility for treatment and examination

**\*\*\*Contact dispatch to notify Police of possible Sexual Assault\*\*\***

# Roane County OES, EMS Division

---

## MEDICAL EMERGENCY

### 315 Sickle Cell Anemia

#### Assessment

History of Sickle Cell Anemia Signs of infection Hypoxia Dehydration Painful joint(s) Limited movement of joints
---------------------------------------------------------------------------------------------------------------------------------

#### EMR

1. Oxygen and airway maintenance appropriate to patient's condition
2. Supportive care
3. **EMR STOP**

#### EMT

4. Pulse oximetry (keep oxygen sats >95%)
5. **EMT STOP**

#### AEMT

6. INT or IV NS TKO
7. **AEMT STOP**

#### PARAMEDIC

8. EKG Monitor, 12 Lead if cardiac concerns present
9. If pain persists:
  - A. **Morphine** 2-5 mg IV/IO or;
  - B. **Fentanyl** 25-50 mcg IV/IO

#### \*\*\*NOTES:

Use caution in administering narcotic to a patient with SpO2 <95%

\*\*\*ALL PATIENTS WHO RECEIVE NARCOTIC MEDICATION MUST BE TRANSPORTED FOR FURTHER EVALUATION

# Roane County OES, EMS Division

## MEDICAL EMERGENCY

### 316 Unconscious / Unresponsive / Altered Mental Status

#### Assessment

Unconscious or unresponsive with vital signs  
Any patient not responding appropriately to verbal or painful stimulus  
Altered level of consciousness with vital signs  
Assess for head trauma  
Assess for Hypo/Hyperthermia, hemiparesis, and fever, OD, Hypoglycemia  
Pediatrics - less commonly associated with intussusception (fold of one intestine into another), intracranial catastrophe, metabolic disorder

#### EMR

1. Oxygen and airway maintenance appropriate for patient's condition
2. Assess for underlying causes: head trauma, hypovolemia, hypothermia, hemiparesis, and fever and treat accordingly
3. **EMR STOP**

#### EMT

4. Pulse oximetry
5. **EMT STOP**

#### AEMT

6. Glucose check
7. IV NS TKO or INT
8. Titrate **Dextrose**, 25 grams IVP (pediatrics 2 cc/kg D<sub>25</sub> IV/IO) through IV with fluids if blood sugar <80 mg/dl.
9. Administer **Narcan** 0.4 mg IV/IM/IN/IO titrated to adequate ventilation (pediatrics <5 y.o - 0.1 mg/kg IV, >5 y.o - 0.4 mg IV). May repeat dose up to 2 mg.
10. **AEMT STOP**

#### PARAMEDIC

11. EKG Monitor, 12 lead if cardiac cause considered
12. Consider **Thiamine** 100 mg/IV if alcoholism or malnourished.
13. If hypotensive, 20 cc/kg NS fluid challenge (pediatrics 20 cc/kg)
14. Contact Medical Control for further orders

**NOTE:** The goal of Narcan therapy is to restore adequate ventilation. Larger doses, especially in patients on chronic opiate therapy, need very small doses of Narcan in the event of overdose. Larger doses of Narcan usually create more agitation and behavioral symptoms.

# Roane County OES, EMS Division

---

## MEDICAL EMERGENCY

### 317 Syncope

#### Assessment

Loss of consciousness with recovery  
Lightheadedness, dizziness  
Palpitations, slow or rapid pulse, irregular pulse Decreased blood pressure

#### EMR

1. Oxygen and airway maintenance appropriate to the patient's condition
2. Supportive Care
3. **EMR STOP**

#### EMT

4. Pulse oximetry
5. **EMT STOP**

#### AEMT

6. Glucose check
7. INT or IV NS TKO - if hypotensive 20 cc/kg bolus (*peds 20 cc/kg bolus*)
8. Titrate **Dextrose** 50% PRN slowly until normal levels achieved, if hypoglycemic
9. **AEMT STOP**

#### PARAMEDIC

10. EKG Monitor, 12 Lead EKG, treat any cardiac dysrhythmia per appropriate protocol
11. Assess neurological status; if abnormal refer to appropriate protocol

# Roane County OES, EMS Division

---

## MEDICAL EMERGENCY

### 318 Pain Management

#### EMR

1. Position as tolerated and supportive care.
2. **EMR STOP**

#### EMT

3. Pulse oximetry
4. **EMT STOP**

#### AEMT

5. INT or IV NS, if hypotensive 20 cc/kg (*pediatrics 20cc/kg*)
6. **AEMT STOP**

#### PARAMEDIC

7. Chest pain - Follow chest pain protocol
8. Abdominal pain - Follow appropriate abdominal pain protocol.
9. Musculoskeletal pain - Follow musculoskeletal pain protocol
10. Kidney Stones - Determine a history. Does the patient have a history of kidney stones?
  - a Do a complete assessment to determine if there are other underlying problems
  - b Utilize appropriate conventions, i.e. O2, IV and EKG monitor
  - c **Toradol** 15 - 30 mg IVP, if no relief from Toradol administer
  - d **Fentanyl** 25 - 50 mcg IVP for severe pain then 50 - 100 mcg every 3 - 5 minutes to a maximum dose of 300 mcg or until relief is achieved
  - e **Morphine** 2 -10 mg IVP up to a maximum 20 mg or until relief is achieved
  - f **Contact Medical Control**
11. Burns - Follow burn protocol

# Roane County OES, EMS Division

---

## SHOCK / TRAUMA

### 401 Air Ambulance Transport

#### PARAMEDIC

Request for an Air Ambulance must be in accordance with approved service policy.

A scene flight by air ambulance MAY be indicated IF:

- The level I trauma patient's condition warrants immediate and extreme action **and** the extrication **and/or** transport time is greater than 30 minutes **and** if the patient is **not** in full trauma arrest.
- Transport time is defined as the length of time beginning when the emergency unit would leave the scene transporting until time of arrival at the trauma center.

The on scene Paramedic or EMS Supervisory Personnel shall have the authority to disregard the response of an air ambulance in accordance with approved service policy.

#### Additional Criteria:

- Multi-system blunt or penetrating trauma with unstable vital signs
- Greater than 25% TBSA burns
- Paralysis or spinal injury
- Amputation proximal to wrist or ankle
- Flail or crushed chest

#### Situational Criteria:

- High energy mechanisms
- Prolonged entrapment
- Multiple casualty incident

Patients will be categorized according to the current Tennessee Trauma Destination Determinates.

- **DO NOT** request an air ambulance transport if patient is in traumatic cardiopulmonary arrest. If the patient has no vital signs, they are in trauma full-arrest.
- The Paramedic in charge of the patient shall have the authority **through** the Incident Commander to disregard the response of the air ambulance.
- The Paramedic will coordinate with the Incident Commander to insure the helicopter receives patient information and landing zone location.

**NOTE:** Medical responsibility will be assumed by the medical flight crew personnel upon arrival at the scene.

**NOTE:** Limitations of the helicopter:

- A. Adults who have traction splint(s) applied
- B. Patients over 6'4"
- C. Patients whose girth exceeds 27"
- D. Any splint or device that exceeds the boundary of the long spine board

## 402 Abdominal/Pelvic Trauma

### Assessment

Abdominal / retroperitoneal abrasions/contusions  
Penetrating injuries  
Hypotension  
Abdominal evisceration(s)  
Abdominal pain on palpation  
Hematuria, bloody stool  
Altered bowel sounds  
Vomiting blood  
History of abdominal injury/trauma  
Suspected injury secondary to mechanism of trauma

### EMR

1. Oxygen and airway maintenance appropriate for the patient's condition
2. C-Spine protection
3. Stop any life threatening hemorrhaging
4. Supportive care
5. **EMR STOP**

### EMT

6. Systolic BP or peds normal for age:
  - A. If Systolic BP >90 mmHg place patient supine with legs elevated and flexed at knees and hips. If no C-Spine concerns, contact Medical Control
7. Patient Pregnant:
  - A. If patient is not past 1<sup>st</sup> trimester: place patient supine with legs elevated and flexed at knees and hips. If no C-Spine concerns, contact Medical Control B. If patient is past 1<sup>st</sup> trimester: place patient in left lateral recumbent position
8. Penetrating object:
  - A. If no penetrating object: place patient supine with legs elevated and flexed at knees and hips. If no C-Spine concerns, contact Medical Control B. If penetrating object present: stabilize object(s)
9. Evisceration:
  - A. If present: place patient supine with legs elevated and flexed at knees and hips. If no C-Spine concerns, contact Medical Control. Cover evisceration(s) with saline soaked trauma dressing
10. Pulse oximetry
11. **EMT STOP**

CONTINUED ON NEXT PAGE

# Roane County OES, EMS Division

---

## SHOCK / TRAUMA

### AEMT

**12. IV NS/LR TKO**

A. If systolic BP <90 mmHg, IV NS/LR 20 cc/kg bolus (*peds 20 cc/kg bolus*).

Target SBP is 90-110 mmHg in adult trauma patients

**13. AEMT STOP**

### PARAMEDIC

**14. EKG Monitor, 12 Lead EKG if cardiac causes a concern**

## 403 Avulsed Teeth - Standing Order

### Assessment

Avulsed teeth may be handled in much the same manner as small parts; i.e. rinse in normal saline (do not rub or scrub) and place in gauze moistened with saline  
Do not cool tooth/teeth with ice

### EMR EMT AEMT

1. Oxygen and airway maintenance appropriate to patient's condition
2. C-Spine stabilization
3. Treat other associated injuries
4. Pay attention to the airway, bleeding and avulsed teeth may cause obstruction.
5. Supportive care
6. Avulsed teeth may be handled in much the same manner as small body parts; i.e. rinse in normal saline (do not rub or scrub) and place in moistened gauze, but there is no need to cool with ice.
7. **EMR, EMT, and AEMT STOP**

### PARAMEDIC

8. Re-implantation at the scene is recommended as this creates maximum possibility of reattachment. The following guidelines pertain to re-implantation at the scene:
  - A. Applicable only for permanent teeth (i.e. with patients generally over 6 years of age)
  - B. Applicable when only one or two teeth are cleanly avulsed and the entire root is present
  - C. Applicable only to anterior teeth (front 6, upper and lower)
  - D. The patient must be conscious
  - E. Should be attempted within the first 30 mins. (The sooner performed the greater the success rate.)
  - F. Do not force re-implantation. Gentle insertion is all that is necessary. Slight incorrect positioning can be corrected later.
9. If re-implantation is not feasible and the patient is a fully conscious adult then the best procedure is to place the tooth in the mouth, either under the tongue or in the buccal vestibule. This is not recommended for children.

## SHOCK / TRAUMA

### 404 Cardiogenic Shock

#### Assessment

Frequently associated with tachy/brady dysrhythmia, acute MI, or blunt chest trauma  
Neck vein distension in sitting position  
Moist sounding lungs (rales, rhonchi)  
Peripheral edema (if chronic heart failure)  
Determine if cardiac dysrhythmia exists  
Consider tension pneumothorax  
Consider cardiac tamponade  
Increased heart rate  
Decreased BP Altered LOC

#### EMR

1. Semi Fowlers or position of comfort
2. Oxygen and airway maintenance appropriate to patient's condition
3. **EMR STOP**

#### EMT

4. Pulse oximetry
5. **EMT STOP**

#### AEMT

6. IV NS or LR, if hypotensive give 20 cc/kg bolus (peds 20 cc/kg bolus)
7. **AEMT STOP**

#### PARAMEDIC

8. EKG Monitor, 12 lead EKG
9. Treat cardiac rhythm appropriately
10. **Contact Medical Control, consider:**
  - A. Dopamine 400 mg / 250 cc or 800 mg / 500 cc D5W IV mix, begin 2-20 ug/kg/min (peds 2-20 ug/kg/min)

## 405 Eye Trauma

### Assessment

Impaled object Inability to open eye(s) Swollen, edematous eye(s) Photophobia Visual defects, loss of vision Redness
-------------------------------------------------------------------------------------------------------------------------------------

<b>EMR EMT AEMT PARAMEDIC</b>
-------------------------------

### Treatment - Standing Order

1. Oxygen and airway maintenance appropriate for patient's condition
2. C-Spine protection if needed
3. If thermal or chemical
  - A. Flush eye(s) with NS or water for 15 min
  - B. Cover both eyes
  - C. Transport
4. Penetration A. Stabilize
  - B. Cover unaffected eye
  - C. Transport
5. Blunt trauma A. Cover both eyes
  - B. Transport
6. Is loss of vision present:
  - A. No - Contact Medical Control
  - B. Yes - If loss of vision was sudden, painless and non-traumatic, consider Retinal Artery Occlusion. Contact Medical Control and:
    - I. Apply cardiac monitor and assess for changes (EMTP only)
    - II. Apply vigorous pressure using heel of hand to affected eye for 3-5 seconds, then release (patient may perform this procedure and may be repeated as necessary)

# Roane County OES, EMS Division

## SHOCK / TRAUMA

### 406 Hypovolemic Shock

#### Assessment

Blood loss due to penetrating injuries to torso or other major vessel  
Fracture of femur or pelvis  
G.I. Bleeding, vaginal bleeding, or ruptured ectopic pregnancy  
Dehydration cause by vomiting, diarrhea, inadequate fluid intake, excessive fluid loss due to fever, uncontrolled diabetes, or burns Pulse may be greater than 120 beats per minute  
Blood pressure may be less than 90 mmHg Systolic  
Orthostatic (Tilt) changes in vital signs (consider possible spinal injury) pulse increase of 20 beats per minute, B decrease of 10 mmHg systolic  
Severe shock (hypovolemia) is defined as decreased level of consciousness, absent radial pulse, capillary refill greater than 2 seconds, no palpable blood pressure

#### EMR

1. Oxygen and airway maintenance appropriate to patient's condition
2. Consider spinal protection
3. Control gross hemorrhage - consider tourniquet or hemorrhage control clamp
4. Trendelenburg patient if no suspected spinal injury
5. **EMR STOP**

#### EMT

6. Pulse oximetry
7. **EMT STOP**

#### AEMT

8. IV NS bolus (20 cc/kg)
9. **AEMT STOP**

#### PARAMEDIC

10. EKG Monitor, 12 Lead EKG
11. IV NS or LR x2 large bore titrated to restore patient's vital signs (in patients with ongoing blood loss maintain patient's systolic blood pressure 90-110 mmHg)
12. Contact Medical Control, Consider: Dopamine 400 mg / 250 ml D5W, 2-20  $\mu$ /kg/min
13. **Pediatrics**
  - A. IV/IO NS 20 cc/kg bolus
  - B. Reassess patient
  - C. Repeat fluid bolus 20 cc/kg if no improvement
  - D. Place a second IV as needed
  - E. Maintain temperature >97°
  - F. Consider Dopamine as above 2-20  $\mu$ /kg/min

## Roane County OES, EMS Division

---

**\*\*\*NOTE:** Cervical spine immobilization is not necessary in patients suffering penetrating trauma (stab or gunshot wound) below the nipple line **AND** no evidence of spinal or head injury. Do not delay transport of patients meeting these criteria for immobilization.

## SHOCK / TRAUMA

### 407 Major Thermal Burn

#### Major Burn:

- Greater than 20% BSA, partial thickness surface involvement
- Greater than 10% BSA, full thickness burn
- Full thickness burns of the head, face, feet, hands or perineum
- Inhalation burn or electrical burns
- Burns complicated by fractures or other significant injury
- Elderly, pediatric, or compromised patients

#### Assessment

Remove clothing from affected parts  
DO NOT pull material out of the burn site: Cut around it  
Look for burns of the nares, oropharangeal mucosa, face or neck  
Listen for abnormal breath sounds  
Note if burn occurred in closed space  
Determine extent of injury (including associated injuries)  
Cardiac monitor for all major burn patients  
Respiratory distress  
ETOH/drug use  
Associated injuries/trauma  
Hypotension  
Past medical history Oropharyngeal burns

#### EMR

1. Stop the burn process with tepid water or normal saline solution and remove any smoldering clothing
2. Oxygen and airway maintenance appropriate to the patient's condition
  - A. Edema may cause patient's airway to close almost instantly without warning signs
  - B. Be prepared to assist ventilation with a BVM
3. Monitor all vital signs and continue reassessment with emphasis on the respiratory rate, peripheral pulses (circulation) and level of consciousness
4. Remove any jewelry
5. Cover burned area with dry sterile dressing or burn sheet. Attempt to keep blisters intact
6. DO NOT use Water-Jel or any other commercially manufactured burn products. DO NOT remove if applied prior to arrival.
7. Monitor to prevent hypothermia
8. Stabilize all associated injuries (e.g. chest, potential spinal injury, fractures, dislocations, etc.)
9. **EMR STOP**

## EMT

10. Pulse oximetry
11. **EMT STOP**

## AEMT

12. INT or IV NS, if hypotensive 20 cc/kg (*peds 20 cc/kg*)
13. **AEMT STOP**

## PARAMEDIC

14. EKG Monitor
15. For major burns, Administer **Morphine** 5-10 mg IV/IO or; **Fentanyl** 25-50 mcg IV/IO (contact medical control in multi-system trauma/pregnancy or for additional pain meds), transport (consider aeromedical transport)
16. If extremity injured, cover open fractures/lacerations/injuries with sterile dressing, splint fractures PRN, avoid unnecessary movement, transport
17. Consider contacting Medical Control for sedating agents especially in pediatric patients
18. Consider cyanide poisoning in obtunded patients and administer Cyanide Antidote if suspected.

Administer IV fluids using the following guide:

- 500 mL per hour for patients over 15 years old
- *250 mL per hour for patients 5 - 15 years old*
- *125 mL per hour for patients under 5 years old*

Excessive or overly aggressive amounts of fluid administration may increase third-spacing shock

# Roane County OES, EMS Division

## SHOCK / TRAUMA

### 408 Musculoskeletal Trauma

#### Assessment

Hypotension  
Past medical history  
Deformity, swelling, tenderness, crepitus, open or closed fractures  
Hemorrhaging, lacerations, ecchymosis, instability  
Decreased function, pulses  
Loss of sensation of distal extremities  
ETOH/drug use  
Mechanism of Injury

#### EMR

1. Oxygen and airway maintenance appropriate for the patient's condition
2. C-Spine protection PRN
3. Control any life threatening hemorrhaging, consider a tourniquet or hemorrhage control clamp
4. **EMR STOP**

#### EMT

5. Consider applying MAST as a splint
6. Splint PRN, stabilize penetrating objects
7. Pulse oximetry
8. **EMT STOP**

#### AEMT

9. INT or IV, LR TKO, if hypotensive 20 cc/kg (*peds 20 cc/kg*)
10. **AEMT STOP**

#### PARAMEDIC

11. EKG Monitor
12. Trauma: Isolated extremity trauma only - consider tourniquet or hemorrhage control clamp. Clamp is approved for scalp use.
  - A. If systolic BP >90 mmHg or peds normal range for age,
    - I. Consider pain management **Morphine** 2-10 mg/IV or **Fentanyl** 25-50 mcg/IV. Can repeat **Fentanyl** to a max dose of 100 mcg.
    - II. Cover open fractures/lacerations, check distal motor/sensory/pulse pre/post splinting, avoid unnecessary movement
  - B. If systolic BP <90 mmHg, IV NS/LR 20 cc/kg (*peds 20 cc/kg*)
  - C. If patient pregnant: Isolated extremity trauma only
    - I. If past the 1<sup>st</sup> trimester and systolic BP >90 mmHg contact Medical Control

## Roane County OES, EMS Division

---

- II. If systolic BP <90 mmHg place patient in left lateral recumbent position, IV NS/LR 20 cc/kg

Cervical spine immobilization is not necessary in patients suffering penetrating trauma (stab or gunshot wound) below the nipple line **AND** no evidence of spinal or head injury. Do not delay transport of patients meeting these criteria for immobilization.

**SHOCK / TRAUMA**

## SHOCK / TRAUMA

### 409 Multi-System Trauma

#### EMR

12. Initiate in-line C-Spine protection while simultaneously evaluating and controlling the patient's ABCs. Incorporate the Mechanism of Injury into the patient care scheme.
13. Control any hemorrhage and simultaneously provide: Oxygen and airway maintenance appropriate to the patient's condition
14. Secure patient to LSB
15. **EMR STOP**

#### EMT

5. Pulse oximetry
6. **EMT STOP**

#### AEMT

7. INT or IV NS, if hypotensive 20 cc/kg (*peds 20 cc/kg*)
8. **AEMT STOP**

#### PARAMEDIC

9. EKG Monitor
10. Consider use of tourniquet or hemorrhage control clamp
11. Refer to specific protocols for specific injuries, consider aeromedical services.
12. Consult medical control for specific orders.

**NOTE: Cervical spine protection is not necessary in patients suffering penetrating trauma (stab or gunshot wound) below the nipple line AND no evidence of spinal or head injury. Do not delay transport of patients meeting these criteria for immobilization.**

## 410 Neurogenic Shock

### Assessment

Associated with spinal cord injuries, closed head injuries and overdoses Signs of hypovolemic shock without pale diaphoretic skin (warm shock)

#### EMR

1. Oxygen and airway maintenance appropriate for patient's condition
2. Establish and maintain C-Spine protection
3. Hemorrhage control
4. Supportive care
5. **EMR STOP**

#### EMT

6. Pulse oximetry
7. **EMT STOP**

#### AEMT

8. INT or IV NS, if hypotensive 20 cc/kg  
*(peds 20 cc/kg)*
9. **AEMT STOP**

#### PARAMEDIC

19. EKG Monitor
20. Contact Medical Control to consider:  
Adult and pediatric - Dopamine at 2-20  $\mu$ /kg/min

**SPECIAL NOTE:** Consider occult bleeding and treat as Hypovolemic Shock Protocol

## SHOCK / TRAUMA

### 411 Septic Shock

#### Assessment

Hot and dry or cool and clammy skin Poor capillary refill Tachycardia/Hypotension Potential for underlying infection
-------------------------------------------------------------------------------------------------------------------------------

#### EMR

1. Oxygen and airway maintenance appropriate for patient's condition
2. Obtain and record oral or axillary temperature if possible
3. **EMR STOP**

#### EMT

4. Pulse oximetry
5. Maintain body temperature above 97° F
6. **EMT STOP**

#### AEMT

7. Glucose check
8. INT or IV NS, if hypotensive 20 cc/kg (*peds 20 cc/kg*)
9. Titrate Dextrose 50% PRN slowly until normal levels achieved, if hypotensive.
10. **AEMT STOP**

#### PARAMEDIC

11. EKG Monitor, 12 Lead EKG if cardiac cause considered
12. If no improvement after two boluses of IV fluids, contact Medical Control and consider:  
Dopamine 2-20  $\mu$ /kg/min (*peds 2-20  $\mu$ /kg/min*)

NOTE: Ensure Body Substance Isolation precautions are utilized.

## SHOCK / TRAUMA

### 412 Soft Tissue / Crush Injuries

#### Assessment

Hypotension  
Past medical history  
Deformity, swelling, tenderness, crepitus, open or closed fractures  
Hemorrhaging, lacerations, ecchymosis, instability  
Decreased function, pulses  
Loss of sensation of distal extremities  
ETOH/drug use  
Mechanism of Injury

#### EMR

1. Oxygen and airway maintenance appropriate for patient's condition
2. C-Spine protection PRN
3. Control any life threatening hemorrhaging
4. **EMR STOP**

#### EMT

5. Consider applying MAST as a splint
6. Other splints PRN, stabilize penetrating objects
7. Pulse oximetry
8. **EMT STOP**

#### AEMT

9. INT or IV, NS LR, if hypotensive 20 cc/kg (*peds 20 cc/kg*)
10. **AEMT STOP**

#### PARAMEDIC

11. EKG Monitor
12. Trauma: Isolated extremity trauma only - consider tourniquet use. iTClamp may be used on scalp lacerations as well.
  - A. If systolic BP >90 mmHg or peds normal range for age,
    - I. Consider pain medications **Morphine** 2-10 mg/IV or **Fentanyl** 25-50 mcg/IV. **Fentanyl** may be repeated to max dose of 100 mcg.
    - II. Cover open fractures/lacerations, check distal motor/sensory/pulse pre/post splinting, avoid unnecessary movement
  - B. If systolic BP <90 mmHg, IV NS LR 20 cc/kg (*peds 20 cc/kg*)
  - C. If patient pregnant: Isolated extremity trauma only

## Roane County OES, EMS Division

---

### SHOCK / TRAUMA

- II. If past 1<sup>st</sup> trimester and systolic BP <90 mmHg contact Medical Control
- III. If systolic BP <90 mmHg place patient in left lateral recumbent position, IV NS LR 20 cc/kg

## SHOCK / TRAUMA

### 413 Spinal Cord Injuries

#### Assessment

Hypotension without actual volume loss Warm/flushed skin despite hypotension Paralysis Loss of reflexes Posturing Priapism Diaphragmatic breathing
----------------------------------------------------------------------------------------------------------------------------------------------------------------------

#### EMR

1. Oxygen and airway maintenance appropriate for the patient's condition
2. C-Spine protection/immobilization
3. Control hemorrhaging

#### EMT

4. Pulse oximetry
5. **EMT STOP**

#### AEMT

6. IV fluid NS LR, if hypotensive bolus 20 cc/kg (repeat bolus once if needed)
7. **AEMT STOP**

#### PARAMEDIC

8. EKG Monitor

SHOCK / TRAUMA

414 Traumatic Cardiac Arrest

Assessment

Cardiac arrest secondary to trauma

**EMR**

1. Oxygen and airway maintenance appropriate for the patient's condition
2. CPR
3. **EMR STOP**

**EMT**

4. Pulse oximetry
5. **EMT STOP**

**AEMT**

6. IV NS/LR give 20 cc/kg bolus
7. Consider second IV access
8. **AEMT STOP**

**PARAMEDIC**

9. EKG Monitor
10. Treat cardiac rhythms per specific protocols
11. If suspected pneumothorax perform needle chest decompression
12. Consider viability of patient prior to transport
13. Contact medical control for instructions or order to cease resuscitation.

## SHOCK / TRAUMA

### 415 Tension Pneumothorax

Patient must meet AT LEAST THREE of the below assessment findings to qualify for this standing order, otherwise, contact Medical Control

#### Assessment

Acute respiratory distress, cyanosis  
Unilaterally decreased breath sounds or absent breath sounds  
Hyper-Resonance of chest unilaterally  
Jugular vein distension  
Subcutaneous Emphysema  
Acute traumatic chest injury, ecchymosis or obvious rib fractures  
History of COPD or other chronic lung disease which predisposes patient to spontaneous pneumothorax  
Hypotension  
Tracheal deviation away from the affected side  
Arrhythmia  
Oxygen saturation - <90% Mechanism of Injury

#### EMR

1. Oxygen and airway maintenance appropriate to patient's condition
2. Perform frequent evaluation of the breath sounds and blood pressure
3. Control any life threatening hemorrhaging
4. **EMR STOP**

#### EMT

5. Consider institution of the multiple trauma protocol, if indicated. Remember this order may be indicated for the medical patient as well.
6. Follow the trauma treatment priority reference as needed
7. If the traumatic tension pneumothorax is secondary to a sucking chest wound, apply an occlusive dressing and treat appropriately
8. Pulse oximetry
9. **EMT STOP**

#### AEMT

10. IV NS LR, If hypotensive 20 cc/kg (peds 20 cc/kg)
11. **AEMT STOP**

SHOCK / TRAUMA

PARAMEDIC

12. EKG Monitor, 12 Lead EKG if cardiac effects present
13. If tension pneumothorax suspected, perform needle decompression per procedure.  
Use 14g 3.5" needle (*pediatrics may use smaller 18g needle*)
14. Contact medical control.

## SHOCK / TRAUMA

### 416 Traumatic Amputation(s)

#### Assessment

Hypotension  
Past medical history  
Deformity, swelling, tenderness, crepitus, open or closed fractures  
Hemorrhaging, lacerations, ecchymosis, instability  
Decreased function, pulses  
Loss of sensation of distal extremities  
ETOH/Drug use  
Mechanism of Injury

#### EMR

1. Oxygen and airway maintenance appropriate for patient's condition
2. C-Spine protection PRN
3. Control any life threatening hemorrhaging
4. **EMR STOP**

#### EMT

5. Consider applying MAST as a splint
6. Other splints PRN
7. Amputated part: If recovered rinse with NS, wrap in moist dressing, place in plastic bag, and transport with patient.
8. Pulse oximetry
9. **EMT STOP**

#### AEMT

10. INT or IV, NS LR, if hypotensive 20 cc/kg (*peds 20 cc/kg*)
11. **AEMT STOP**

#### PARAMEDIC

12. EKG Monitor
13. Amputation - consider tourniquet use
14. If systolic BP >90 mmHg or peds normal range for age consider **Morphine** 2-10 mg/IV or **Fentanyl** 25-50 mcg/IV. **Fentanyl** may be repeated to max 100 mcg.
15. Cover open fractures/lacerations, check distal motor/sensory/pulse pre/post splinting, avoid unnecessary movement
16. Contact medical control.

## OBSTETRICAL EMERGENCIES

### 500 Obstetrical / Gynecological Complaints (Non-Delivery or GYN only)

#### Assessment

Patient Para (number of live births) and Gravida (number of pregnancies)  
Term of pregnancy in weeks, EDC, Multiple births expected or history  
Vaginal bleeding (how long and approximate amount)  
Possible miscarriage/products of conception  
Pre-natal medications, problems, and care  
Last menstrual cycle  
Any trauma prior to onset?  
Lower extremity edema

#### EMR

1. Oxygen and airway maintenance appropriate for the patient's condition
2. Patient positioning appropriate for condition
3. **EMR STOP**

#### EMT

4. Control hemorrhage as appropriate
5. Pulse oximetry
6. **EMT STOP**

#### AEMT

7. Glucose check
8. INT or IV NS TKO unless signs of shock, then 20 cc/kg fluid bolus
9. **AEMT STOP**

#### PARAMEDIC

10. EKG Monitor
11. Contact medical control.

## OBSTETRICAL EMERGENCIES

### 501 Normal Delivery

#### Assessment

Patient Para (number of live births) and Gravida (number of pregnancies) Term of pregnancy in weeks, EDC Vaginal Bleeding Pre-natal medications, problems, and care Membrane ruptured Lower extremity edema
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

#### EMR

#### Mother:

1. Oxygen and airway maintenance appropriate for patient's condition
2. **EMR STOP**

#### EMT

#### Mother:

3. Pulse oximetry
4. Check mother for crowning, PRN
5. Use gentle pressure to control delivery. When head delivers, suction airway and check for cord around neck
6. After delivery, keep mother and infant on same level, clamp cord @ 8 and 10 inches from the baby and cut between clamps
7. Dry infant and wrap to keep warm. Maintain airway
8. Check APGAR at 1 and 5 minutes post-delivery
9. DO NOT allow mother to nurse until both have been evaluated in the Emergency Department
10. Allow placenta to deliver
  - A. Massage uterine fundus (lower abdomen)
  - B. Observe and treat signs of shock with increased delivery of oxygen and IV fluids
  - C. Be alert to the possibility of multiple births
11. Re-Evaluate vaginal bleeding

#### Infant:

1. Protect against explosive delivery
2. When head delivers suction airway (mouth first then nose) & check for cord around neck
3. After delivery clamp cord @ 8 and 10 inches from baby and cut between clamps
4. Dry infant and wrap to keep warm (silver swaddler). Maintain airway, suction PRN
5. Check APGAR Score at 1 and 5 minutes after delivery
6. DO NOT allow infant to nurse until both have been evaluated in the Emergency Department
7. Re-Evaluate cord for bleeding, if bleeding add additional clamp and re-evaluate.

# Roane County OES, EMS Division

## 8. EMT STOP

### AEMT

9. INT or IV LR TKO , if patient in active labor defined as: regular contractions q 3-5 mins with 30-60 second duration.

(Cont. on next page)

### PARAMEDIC

10. EKG monitor

#### NOTE: Considerations

1. The greatest risks to the newborn infant are airway obstruction and hypothermia. Keep the infant warm (silver swaddler), dry, covered, and the infant's airway maintained with bulb syringe. Always remember to squeeze the bulb prior to insertion into the infant's mouth or nose.
2. The greatest risk to the mother is post-partum hemorrhage. Watch closely for signs of hypovolemic shock and excessive vaginal bleeding.
3. Spontaneous or induced abortions may result in copious vaginal bleeding. Reassure the mother, elevate legs, treat for shock, and transport.
4. Record a blood pressure and the presence or absence of edema in every pregnant woman you examine, regardless of chief complaint

Complete patient care reports on BOTH mother and child.

## APGAR SCORING CHART

\*\*\*SCORES OF LESS THAN 6 REQUIRE RESUSCITATIVE MEASURES.

<u>Clinical Sign</u>	<u>0 Points</u>	<u>1 Point</u>	<u>2 Points</u>
Appearance	Blue/Pale	Body Pink Extremities Blue	Completely Pink
Pulse	Absent	Below 100/minute	Above 100/minute
Grimace	No response	Grimace	Cries
Activity	Limp	Some flexion of extremities	Action motion
Respiratory	Absent	Slow/Irregular	Good strong cry

The APGAR score should be calculated after birth of the infant. The five (5) clinical signs are evaluated according to the scoring system detailed above. Each sign is assigned points to be

## Roane County OES, EMS Division

---

totalled. A total score of 10 indicates that the infant is in the best possible condition. A score of 4 to 6 indicates moderate depression and a need for resuscitative measures.

**DO NOT** delay resuscitation efforts to obtain APGAR score. Obtain APGAR at 1 and 5 min.

## OBSTETRICAL EMERGENCIES

### 502 Abruptio Placenta

#### Assessment

Multiparity Maternal hypertension Trauma Drug use Increased maternal age History Vaginal bleeding with no increase in pain No bleeding with low abdominal pain
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

#### EMR

1. Oxygen and airway maintenance appropriate to the patient's condition
2. Position patient in the left lateral recumbent position
3. **EMR STOP**

#### EMT

4. Pulse oximetry
5. **EMT STOP**

#### AEMT

6. IV NS TKO, if hypotensive 20 cc/kg
7. **AEMT STOP**

#### PARAMEDIC

8. EKG Monitor

## OBSTETRICAL EMERGENCIES

### 503 Amniotic Sac Presentation

#### Assessment

Amniotic sac visible  
Membrane not broken  
Fetus may or may not be visible  
Pre-natal medications, problems, and care  
Usually third trimester  
Applies to greater than 20 weeks gestation  
Abdominal pain  
Indications of immediate delivery

#### EMR

1. Oxygen and airway maintenance appropriate to the patient's condition
2. Place patient in a position of comfort
3. **EMR STOP**

#### EMT

4. Amniotic sac  
If no fetus visible, cover presenting part with moist, sterile dressing  
If head of the fetus has delivered, tear sac with fingers and continue steps for delivery
5. Contact Medical Control ASAP
6. Pulse oximetry
7. **EMT STOP**

#### AEMT

8. IV NS TKO, if hypotensive 20 cc/kg (peds20 cc/kg)
9. **AEMT STOP**

#### PARAMEDIC

10. EKG Monitor

## OBSTETRICAL EMERGENCIES

### 504 Breech or Limb Presentation

#### Assessment

Patient para (number of live births) and gravida (number of pregnancies)  
Term of pregnancy in weeks, EDC  
Vaginal bleeding  
Pre-natal medications, problems, and care  
Water broken  
Buttock, arm or leg presentation

#### EMR

1. Oxygen and airway maintenance appropriate to patient's condition
2. **EMR STOP**

#### EMT

3. Pulse oximetry

#### Breech Presentation - Treatment - Standing Order - All EMTs 4.

Allow the delivery to progress spontaneously - DO NOT PULL!

5. Support the infant's body as it delivers
6. If the head delivers spontaneously, deliver the infant as noted in 'Normal Delivery'
7. If the head does not deliver within 3 minutes, insert a gloved hand into the vagina and maintain airway for the infant
8. DO NOT remove your hand until relieved by a Higher Medical Authority.

#### Limb Presentation - Treatment - Standing Order - All EMTs

9. Position the mother in a supine position with the head lowered and pelvis elevated
10. **EMT STOP**

#### AEMT

11. IV NS TKO, if hypotensive 20 cc/kg (*peds 20 cc/kg*)
12. **AEMT STOP**

#### PARAMEDIC

13. EKG Monitor
14. Transport ASAP

## OBSTETRICAL EMERGENCIES

### 505 Meconium Stain

#### Assessment

Patient para (number of live births) and gravida (number of pregnancies)  
Term of pregnancy in weeks, EDC  
Vaginal bleeding  
Pre-natal medications, problems, and care  
Membrane ruptured  
Amniotic fluid that is greenish or brownish yellow Fecal material expelled with the amniotic fluid

#### EMR

1. Do not stimulate respiratory effort before suctioning the oropharynx
2. Suction the **mouth then the nose** (using a meconium aspirator) while simultaneously providing Oxygen 100% by blow by method and while maintaining the airway appropriate to the patient's condition
3. **EMR STOP**

#### EMT

4. Pulse oximetry
5. Obtain and APGAR score after airway treatment priorities. Score one minute after delivery and at five minutes after delivery. (Time permitting)
6. Repeat initial assessment and complete vital signs until patient care is transferred to the appropriate ER staff.
7. **EMT STOP**

#### AEMT

8. IV NS TKO, if hypotensive 20 cc/kg  
*(peds 20 cc/kg)*
9. **AEMT STOP**

#### PARAMEDIC

10. EKG Monitor
11. Contact medical control

## OBSTETRICAL EMERGENCIES

### 506 Placenta Previa

#### Assessment

Painless bleeding which may occur as spotting or recurrent hemorrhage  
Bright red vaginal bleeding usually after 7<sup>th</sup> month  
History  
Multiparity  
Increased maternal age  
Recent sexual intercourse or vaginal exam  
Patient para (number of live births) and gravida (number of pregnancies)  
Term of pregnancy in weeks  
Pre-natal medications, problems, and care  
History of bed rest  
Placenta protruding through the vagina

#### EMR

1. Oxygen and airway maintenance appropriate to patient's condition
2. Position of comfort
3. **EMR STOP**

#### EMT

4. Pulse oximetry
5. **EMT STOP**

#### AEMT

6. IV NS TKO, if hypotensive 20 cc/kg  
*(peds 20 cc/kg)*
7. **AEMT STOP**

#### PARAMEDIC

8. EKG Monitor

**NOTE:** Any painless bleeding in the last trimester should be considered Placenta Previa until proven otherwise. If there are signs of eminent delivery membrane rupture is indicated followed by delivery of the baby. The diagnosis of eminent delivery depends on the visual presence of the baby's body part through the membrane.

**OBSTETRICAL EMERGENCIES**

**507 Prolapsed Umbilical Cord**

Assessment

Cord emerges from the uterus ahead of baby  
With each uterine contraction the cord is compressed between the presenting part and the pelvis  
Pulse on exposed cord may or may not be palpable  
Patient Para (number of live births) and Gravida (number of pregnancies)  
Term of pregnancy in weeks, EDC  
Vaginal bleeding  
Pre-natal medications, problems, and care  
Membrane ruptured

**EMR**

1. Oxygen and airway maintenance appropriate for the patient's condition
2. **EMR STOP**

**EMT**

3. Palpate pulses in the cord
4. Pulse oximetry
5. Position the mother with hips elevated
  - A. Knee to chest
  - B. Hips elevated as much as possible on pillows
6. Instruct mother to pant with each contraction, which prevents her from bearing down.
7. Check for a pulse in the cord
  - A. If no pulse - insert a gloved hand into the vagina and gently push the infant's head off the cord. While pressure is maintained on the head cover the exposed cord with a sterile dressing moistened in saline. Transport immediately and **DO NOT** remove your hand until relieved by hospital staff.
  - B. If pulse present - cover exposed cord with moist dressing
8. Contact Medical Control as soon as possible if time and patient condition allows
9. **EMT STOP**

**AEMT**

10. IV NS TKO, if hypotensive 20 cc/kg (*peds 20 cc/kg*)
11. **AEMT STOP**

**PARAMEDIC**

12. EKG Monitor

## OBSTETRICAL EMERGENCIES

### 508 Pre-eclampsia and Eclampsia

#### Assessment

Patient Para (number of live births) and Gravida (number of pregnancies)  
Term of pregnancy in weeks, EDC  
Vaginal bleeding  
Pre-natal medications, problems, and care  
Membrane ruptured  
Usually begins after the twentieth week of pregnancy  
Most often affects women during their first pregnancy  
May have a history of chronic hypertension and/or diabetes  
May experience hypertension and edema  
May experience headaches, blurred vision, and abdominal pain  
May experience seizures which indicates a progression from pre-eclampsia to eclampsia

#### EMR

1. Oxygen and airway maintenance appropriate for the patient's condition
2. Place patient in left lateral recumbent position
3. **EMR STOP**

#### EMT

4. Pulse oximetry
5. **EMT STOP**

#### AEMT

6. Glucose check
7. IV NS TKO, if hypotensive 20 cc/kg (*peds 20 cc/kg*)
8. **AEMT STOP**

#### PARAMEDIC

9. EKG monitor
10. Versed 2-5 mg IVP per seizure protocol if generalized seizure activity or, if no IV, Versed 10 mg IM (>39 kg) or 5 mg IM (<39 kg)
11. Contact Medical Control and consider:  
Magnesium Sulfate 1-2 grams IV Slowly

**NOTE:** Record a blood pressure and the presence or absence of edema in every pregnant woman you examine no matter what the chief complaint

## MISCELLANEOUS

### 601 Discontinuation / Withholding of Life Support

Once life support has been initiated in the field, **Non ALS personnel CAN NOT** discontinue resuscitative measures unless directed to do so by the on-scene physician, EMT-Paramedic or presented with a valid Physician Orders for Scope of Treatment (POST/DNR).

#### Withholding Resuscitation - Standing Orders

If there is no CPR in progress, CPR may be withheld if one or more of the following conditions are met:

- A. Obviously dead patients with dependent lividity, rigor mortis, or massive trauma (i.e., evacuation of the cranial vault, crushed chest, crushed head, etc.)
- B. Obviously dead patients with tissue decomposition
- C. Patients without vital signs who cannot be accessed for treatment due to entrapment for prolonged time. (12-15 minutes or greater)
- D. Severe blunt trauma with absence of BP, pulse, respiratory effort, neurologic response, and pupillary response
- E. When presented a valid POST/DNR order or a copy as approved by the Tennessee Department of Health. DNR and POST orders not on the official state form can be accepted if it is documented in a medical record such as a nursing chart, hospice care, or home nursing
- F. Instructed to do so by the on scene Paramedic

#### Discontinuing Life Support

Once life support has been initiated in the field, in order to discontinue life support, the following conditions must be met:

1. Asystole is present on the EKG monitor in two leads **and**
2. There is an absence of pulse, respirations, and neurological reflexes **and**
3. At least one of the following conditions are met:
  - A. Appropriate airway management has been confirmed, the patient has been well ventilated with 100% oxygen and multiple (at least three) doses of Epinephrine and have not been effective in generating an EKG complex
  - B. Transcutaneous pacing, if available, has not been effective in generating a pulse
  - C. Obvious signs of death in the absence of hypothermia, cold water drowning, or induced coma, **or**
  - D. The Paramedic can document lack of CPR for at least 10 minutes, **or**
  - E. Prolonged resuscitation (25 minutes of resuscitation with agonal or asystolic rhythm) in the field without hope for survival, **or**
  - F. Massive trauma such as evacuation of cranial vault, etc., **or**
  - G. Sever blunt trauma with absence of vital signs and pupillary response
  - H. End tidal CO<sub>2</sub> less than 10 while performing effective CPR

Upon termination in the field any tubes, needles and IV lines will be left in place (IV lines to be tied off and cut with catheter left in place).

### MISCELLANEOUS

#### NOTES:

- Personnel shall give careful consideration when using this standing order. Conditions such as: overdose, electrical shock, hypothermia, and hypoglycemia may mimic some of the above signs and symptoms.

All deaths must be confirmed by a Paramedic

## MISCELLANEOUS

### 602 Field Determination of Death

#### Assessment

Pulseless, non-breathing with definitive signs of death:  
Rigor Mortis  
Dependent lividity  
Decomposition of body tissue  
Devastating, un-survivable injury  
Decapitation  
Incineration  
Separation of vital internal organ(s) from the body or total destruction of organs  
Gunshot wound to the head that crosses the midline (entrance and exit)

**EMR EMT AEMT PARAMEDIC**

If patient is pulseless, non-breathing without definitive signs of death:  
Must receive resuscitation unless a properly executed DNR or POST form is present

#### Treatment - Standing Order

##### **DNR Orders:**

- If family member or caregiver can produce a properly executed DNR or POST order, resuscitation can be withheld.
- Treat patients with known DNR orders appropriately; just do not initiate CPR if they develop cardiovascular or respiratory arrest.
- When there is any doubt about what to do, begin resuscitative efforts with all skill available.

##### **Resuscitation has been initiated prior to EMS arrival:**

Anytime CPR or an attempt at resuscitation has been initiated by anyone at the scene, resuscitative efforts will be continued until:

- Medical Control directs the team to stop (either on line or on-scene)
- It is determined the patient meets criteria for “definitive signs” of death

A properly executed DNR or POST form is presented

**MISCELLANEOUS**

**603 Mandatory EKG**

**PARAMEDIC**

EKG monitoring will be mandatory under the following conditions:

**Patients**

- Complaining of chest pain regardless of source
- In cardiac arrest with or without CPR in progress
- That are non-viable (other than those exhibiting body decomposition, dependent lividity, rigor mortis, decapitation)

EKGs will have the following information printed on the recording:

- Name or report number
- Age (if possible)
- Unit number and date

EKGs will be appended to the PCR appropriately

12 Lead EKGs may be applied by any provider EMT or higher on scene; however treatment decisions may only be made by a paramedic.

## MISCELLANEOUS

### 604 Patient Refusal or Declination of Care / Patient Non-Transport Situations

#### Assessment

Determine presence of injury or illness and desire for transport Identify the person who made the EMS call Reason for refusal
-------------------------------------------------------------------------------------------------------------------------------------

<b>EMR EMT AEMT PARAMEDIC</b>
-------------------------------

#### Standing Orders

1. Perform and document mini-mental status exam to confirm mental capacity to refuse care
2. Confirm and document the absence of intoxicating substance or injury
3. Confirm patient is of legal age of majority, or emancipated minor
4. Document mechanism of injury or circumstances of illness
5. Document pertinent past history
6. Perform vital signs and problem directed exam **The following may not refuse transport:**
  1. Patients with impaired judgment and decreased mental status (Utilize the mini mental status exam on next page to determine; document)
  2. Minors (less than 18 years of age or older unless they are emancipated by the courts)
  3. All minors must have refusal from parent or guardian, not older sibling or other relative, unless every effort has been made to contact parent/guardian and was not successful
  4. Do not release minor on the scene without parent/guardian consent

#### Reasons for Non-Transport

Minor illness or injury and acceptable alternative transportation available

#### No Patient Found on the Scene

Definition: No person found to have any complaint of injury/illness of any type or degree PCR is to be completed in detail as to why no patient was found, i.e.: no person found on scene, person located with no complaint of injury/illness and denies needing medical assistance.

## Roane County OES, EMS Division

---

### Mini Mental Status Exam

1. Orientation to time - time of day, day, week, month, year	5 pts max
2. Orientation to place - building, street, city, state, country	5 pts max
3. Say "boy, dog, ball" and have the patient repeat it	3 pts max
4. Ask the patient to spell would backward, or do serial 3s backward from 20	5 pts max
5. Without repeating the words, ask them to repeat the previous three words (boy, dog, ball)	3 pts max
6. Ask the patient to do the following after you have completed the request "stick out your tongue and touch your right hand to your left ear"	3 pts max
7. Ask the patient to identify your pen and watch	2 pts max
8. Ask the patient to read the following sentence then do as it says "Shut your eyes"	1 pt
9. Ask the patient to write a sentence	1 pt
10. Ask the patient to draw two overlapping pentagons (show them an example)	1 pt

A score of 21 or better is considered mentally competent by most psychiatrists for a patient to make reasonable decisions.

## MISCELLANEOUS

### 605 Physical Restraint

**EMR EMT**

#### All Patients:

1. Safety of EMS personnel is the main priority in any situation where a patient exhibits aggressive or combative behaviors and needs to be restrained.
2. Use the minimum amount of force and restraint necessary to safely accomplish patient care and transportation with regard to the patient's dignity. Avoid unnecessary force.
3. Assure that adequate personnel are present and that police assistance has arrived, if available, before attempts to restrain patient.
4. Plan your approach and activities before restraining the patient.
5. Have one person talk to and reassure the patient throughout the restraining procedure.
6. Approach with a minimum of four persons, one assigned to each limb, all to act at the same time.
7. Initial take down may best be accomplished leaving the patient in the prone position. After restraint, the patient should be placed in a supine position.
8. Call for additional help if patient continues to struggle against restraint.
9. Restrain all 4 extremities with patient supine on stretcher.
10. Use soft restraints to prevent the patient from injuring him or herself or others.
11. A police officer or other law enforcement personnel shall always accompany a patient in the ambulance if the patient has been restrained.
12. Do not place restraints in a manner that may interfere with evaluation and treatment of the patient or in any way that may compromise patient's respiratory effort.
13. Evaluate circulation to the extremities frequently.
14. Thoroughly document reasons for restraining the patient, the restraint method used, and results of frequent reassessment.
15. Initial "take down" may be done in a prone position to decrease the patient's visual field and stimulation, and the ability to bite, punch, and kick. After the individual is controlled, he/she shall be restrained to the stretcher or other transport device in the supine position.
16. **DO NOT** restrain patient in a hobbled, hog-tied, or prone position.
17. **DO NOT** sandwich patient between devices, such as long boards or Reeve's stretchers, for transport. Devices like backboards should be padded appropriately.
18. A stretcher strap that fits snugly just above the knees is effective in decreasing the patient's ability to kick.
19. Padded or leather wrist or ankle straps are appropriate. Handcuffs and plastic ties are not considered soft restraints.
20. Never apply restraints near the patient's neck or apply restraints or pressure in a fashion that restricts the patient's respiratory effort.

**(Continued on Next Page)**

## Roane County OES, EMS Division

---

21. Never cover a patient's mouth or nose except with a surgical mask or a NRB mask with high flow oxygen. A NRB mask with high flow oxygen may be used to prevent spitting in a patient that also may have hypoxia or another medical condition causing his/her agitation, but a NRB should never be used to prevent spitting without also administering high flow oxygen through the mask.

### Performance Parameters

1. Verbal techniques include:
  - A. Direct empathetic and calm voice.
  - B. Present clear limits and options.
  - C. Respect personal space.
  - D. Avoid direct eye contact.
  - E. Non-confrontational posture.
2. There is a risk of serious complications or death if patient continues to struggle violently against restraints. Chemical restraint by sedation may be indicated in some dangerous, agitated patients.

### AEMT

3. INT or IV NS/LR, if hypotensive 20 cc/kg (peds 20 cc/kg)
4. **AEMT STOP**

### PARAMEDIC

5. Administer the following if needed and if available in the service formulary:
  - A. Haldol 5 mg IM, repeat once
  - B. Versed: 2-5 mg IV or 5-10 mg IM, repeat once

### Documentation:

Review for documentation of frequent reassessment of vital signs, cardiopulmonary status, and neurovascular status or restrained extremities, reason for restraint and method used. Benchmark of documenting of these items is at least every 15 minutes.

## MISCELLANEOUS

### 606 Physician On Scene

#### EMR, EMT, AEMT, PARAMEDIC

If private physician intervenes by phone the EMT-Basic/EMT-Paramedic shall:

- Request the physician contact Medical Control and relay any orders through them.
- **NO ORDERS** will be taken over the phone from the private physician.

#### Standing Order:

1. No one will be recognized as a physician without proof of license. This must be in the form of a wallet card or visual personal recognition. NO ORDERS will be accepted until proof of license is verified.
2. Consider need for Law Enforcement if any difficulty with person occurs.
3. The EMT or above shall:
  - a. Inform the physician that they must contact Medical/Trauma Control.
  - b. Inform Medical/Trauma Control of the presence of a physician on scene.
4. Medical/Trauma Control may:
  - a. Speak to the physician to determine the qualifications.
  - b. Request the EMT, AEMT, or Paramedic to verify licensure of the physician.
  - c. Relinquish total responsibility for the patient to the on-scene physician.
5. Physician (intervening) may:
  - a. Assist the EMT, AEMT, or Paramedic and allow you to operate under EMS standing orders and protocols. Offer assistance by allowing the EMS Provider to remain under Medical/Trauma Control; or
  - b. Request to talk to Medical/Trauma Control to offer advice and assistance; or
  - c. Take responsibility for the care given by the EMS Provider if okay with Medical/Trauma Control, then physically accompany the patient to the Emergency Department where responsibility is assumed by the receiving physician; and shall,
  - d. Sign for all instructions given to the EMS Provider
  - e. Contact should be made with Medical/Trauma Control if this happens.
6. If private physician intervenes by phone or in person the EMS provider shall:
  - a. Inform the physician that the EMS Provider must contact Medical/Trauma Control.
  - b. Request the physician contact Medical Control and relay any orders through them.

NO ORDERS should be taken over the phone from the private physician. At no time should any order be taken over the telephone except from Medical/Trauma Control.

**MISCELLANEOUS**

**607 By-standers On Scene**

**EMR, EMT, AEMT, PARAMEDIC**

Standing Order:

By-stander participation - You may use them at your discretion. However YOU will be responsible for their actions and treatment. This includes other medical professionals. In any situation you need assistance you may utilize their expertise and skills.

**NOTE:** Request proof of their licensure by visualization of their current license, if possible. Remember, YOU are responsible for the patient. If any by-stander is trying to take over direction of patient care, other than a Physician (follow Guideline 606 Physician on Scene in this situation) you may have law enforcement remove the person for “Obstruction of Emergency Services”.

## MISCELLANEOUS

### 608 Procedure for Deviation from Standing Orders

#### EMR, EMT, AEMT, PARAMEDIC

NEVER simply disregard a standing order or protocol.

These Standing Orders have been established so that EMS Personnel may provide the best care possible for our patients. Most of our patients will be covered by a single Standing Order. However, some patients may have signs and symptoms of illness and/or injury that are covered by more than one Standing Order or, in rare cases, following a Standing Order may not be in the best interest of the patient. In these cases you must be aware that combining Standing Orders may lead to medication errors, overdose, and medication incompatibility. You are expected to use your judgment and to always make decisions that are in the best interest of the patient.

If you use more than one standing order when treating your patient, you must document your reasoning in the NARRATIVE SECTION of the Patient Care Report.

If in your judgment, following a standing order is not in the best interest of the patient, CONTACT MEDICAL CONTROL, regarding your treatment. Document the rationale for deviation, and the name of the physician giving the order.

## MISCELLANEOUS

## 609 Spinal Immobilization

### EMR, EMT, AEMT, PARAMEDIC

The intent of this guideline is to decrease injury and discomfort to patients caused by unnecessary spinal immobilization and use of long spine boards.

- Studies show that immobilizing trauma victims may cause more harm than good to the patient.
- Penetrating trauma victims benefit the most from rapid assessment and transport to a trauma center without spinal immobilization/**spinal motion restriction (SMR)**.
- There is evidence that backboards result in harm by causing pain, changing the normal anatomic lordosis of the spine, inducing patient agitation, causing pressure ulcers, and compromising respiratory function
- Backboards should be avoided for spinal immobilization with conscious patients
- **Placing ambulatory patients on backboards is unacceptable**
- **Use of the backboard is recommended in the event of CPR**

#### Spinal Injury Assessment Introduction:

- Omit SMR if all assessment criteria are safely assessed and normal
- Consider SMR for a patient who is suspected of having a traumatic unstable spinal column injury. Have a high index of suspicion for pediatrics and patients with degenerative skeletal/connective tissue disorders (i.e. osteoporosis, elderly, previous spinal fractures, etc.)
- Penetrating trauma such as a gunshot wound or stab wound should **NOT** be immobilized on a long board unless there are signs of spinal injury. Emphasis should be on airway and breathing management, treatment of shock, and rapid transport to a Level 1 trauma center.
- Determination that immobilization devices should be used or removed should be made by the highest level provider on scene.
- If the immobilization process is initiated prior to the arrival and assessment by the highest level of provider, **STOP** and perform spine injury assessment to determine the best course of action.

#### Pediatric Patients and Car Seats

- **Infants restrained in a rear-facing car seat** may receive SMR and be extricated in the car seat. The child may remain in the seat if the SMR is secure and his/her condition allows (no signs of respiratory distress or shock)
- **Children restrained in a car seat (with a high back - convertible or booster)** may receive SMR and be extricated in the car seat. The child may remain in the seat if the SMR is secure and his/her condition allows (no signs of respiratory distress or shock).
- **Children restrained in booster seat (without a back)** need to be extricated and receive standard SMR procedures.

## Helmet Removal

Safe and proper removal of the helmet should be done following the steps outlined in an approved trauma curriculum.

Indications for football helmet removal:

- When a patient is wearing a helmet and not shoulder pads
- In the presence of head and/or facial trauma, and removal of face piece is not sufficient
- Patients requiring advanced airway management when removal of the facemask is not sufficient
- When the helmet is loose on the patient's head
- In the presence of cardiopulmonary arrest. (The shoulder pads must also be removed.)

When helmet and shoulder pads are both on the spine is kept in neutral alignment. If the patient is wearing only a helmet or shoulder pads, neutral alignment must be maintained. Either remove the other piece of equipment or pad under the missing piece. *All other helmets must be removed in order to maintain spinal alignment.*

## Spinal Motion Restriction

The term spinal motion restriction (SMR) better describes the procedure used to care for patients with possible unstable spinal injuries. SMR includes:

- Reduction of gross movement by patient
- Prevention of duplicating the damaging mechanism to spine
- Regular reassessment of motor/sensory function

### Purpose:

To decrease the risk of negative effects cause by traditional spinal immobilization still providing appropriate care to patients with possible spinal injury by implementing alternative methods to achieve SMR.

### Indications:

Any patient identified whose assessment warrants spinal motion restriction. The spinal injury assessment should be performed prior to application of SMR.

### Procedure:

If patient experiences negative effects of SMR methods used, alternative methods should be utilized.

- A. If hard backboard utilized for extrication, patient should be removed from the extrication device as soon as possible and placed on the ambulance stretcher.
- B. Patients with potential c-spine or spinal column injury should be transported supine directly on flat cot **without** a long spine board. If patient was extricated to stretcher on a long spine board (LSB), unstrap and log-roll the patient, remove the long spine board for transport, and transport on cot.
- C. May be left on LSB if not for spinal immobilization (e.g. extremity splinting) or removal would delay transport of an unstable patient.
- D. Patient positions and/or methods/tools to achieve SMR that are allowable (less invasive to more invasive)

## Roane County OES, EMS Division

---

- E. Patient position: supine, lateral, semi fowlers, fowlers
- F. Tools/methods to achieve position of comfort include, but not limited to: pillows, children's car seat, scoop, vacuum mattress  
\*\*\* (Continued on next page)
- G. Provide manual stabilization restricting gross motion. Alert and cooperative patients may be allowed to self-limit motion if appropriate with or without cervical collar.
- H. Apply cervical collar; patients who are unable to tolerate cervical collar may benefit from soft collars, pillows, or other padding.
- I. Considerations for patient movement when decision to SMR has been made:
- J. Keeping with the goal of restricting gross movement of spine and preventing increased pain and discomfort, self-extrication of the patient is allowable.
- K. If needed, extricate patient limiting flexion, extension, rotation and distraction of spine
- L. Pull sheets, other flexible devices, scoops, and scoop like devices can be employed if necessary. Hard backboards should only have limited utilization.
- M. **No standing take downs of ambulatory patients.** Ambulatory patients who meet the above criteria for cervical immobilization should have c-collar applied and be allowed to sit onto the stretcher.
- N. Apply adequate padding to prevent tissue ischemia and increase comfort. **Patients should be allowed to be in a position of comfort.**
- O. Place patient in position best suited to protect airway
- P. Regularly reassess motor/sensory function (include finger abduction, wrist/finger extension, plantar/dorsal flexion, and sharp/dull exam if possible.
- Q. Consider the use of SpO<sub>2</sub> and EtCO<sub>2</sub> to monitor respiratory function.
- R. Delivery to hospital: movement of patient to hospital stretchers should be done by limiting motion of the spine.

### Special Considerations:

- **Patients with acute or chronic difficulty breathing:** SMR has been found to limit respiratory function an average of 17% with the greatest effect experienced by geriatric and pediatric subjects restricted to a hard backboard. **USE SMR WITH CAUTION with patients presenting with dyspnea and position appropriately.**
- **Pediatric patients, < 9 years of age:**
  - o Consider use of padded pediatric motion restricting board o
  - o Avoid methods that provoke increased spinal movement
  - o If choosing to apply SMR to patient in car seat, ensure that proper assessment of patient posterior is performed

# Roane County OES, EMS Division

- **Combative patients:** Avoid methods that provoke increased spinal movement and/or combativeness
- **Interfacility transports:** Long spine boards do not have a role for patients being transported between facilities. If the sending facility is asking EMS to use a LSB for transport, EMS providers should discuss NOT using a LSB with the sending facility physician before transporting a patient.

## Protocol Guidelines

### *Selective Spinal Immobilization C-Spine Clearance*

Neuro Exam: Any focal deficit?

YES

NO

**Spinal Immobilization Not Required**

NO

YES

Spinal Immobilization Guidelines:  
Spinal Immobilization Procedure

Age 65 or greater / 5 or less and Significant mechanism of Injury?

NO

YES

Alertness: Alteration in mental

NO

YES

Intoxication: Any evidence?

NO

YES

Distracting Injury: Any painful injury that might distract the patient from the pain of c-spine injury?

NO

YES

Spinal Exam: Point tenderness over the spinous process(es) or pain to

Spinal Immobilization Procedure

YES  
Immobilization Procedure

Spinal Immobilization Procedure

Spinal Immobilization Procedure

**Any doubt always immobilize**

1. Long spine boards (LSB) have both risks and benefits for patients and have not been shown to improve outcomes. The best use of the LSB may be for extricating the unconscious patient, or providing a firm surface for compressions. However, several devices may be appropriate for patient extrication and movement, including the scoop stretcher and soft body splints.
2. Utilization of the LSB should occur in consideration of the individual patient's benefit vs. risk.

## Roane County OES, EMS Division

---

3. Patients who should be immobilized with a LSB include: Patients with blunt trauma and distracting injury, intoxication, altered mental status, or neurologic complaint (e.g. numbness or weakness), and non-ambulatory blunt trauma patients with spinal pain, tenderness, or spinal deformity.
4. Patients with penetrating trauma and no evidence of spinal injury do not require spinal immobilization. Patients who are ambulatory at the scene of blunt trauma in general do not require immobilization via LSB, but may require cervical collar and spinal precautions.
5. Whether or not a LSB is utilized, spinal precautions are **STILL VERY IMPORTANT** in patients at risk for spinal injury. Adequate spinal precautions may be achieved by placement of a hard cervical collar and ensuring that the patient is secured tightly to the stretcher, ensuring minimal movement and patient transfers, and manual in-line stabilization during any transfers.

## MISCELLANEOUS

### 610 Stretcher Transport

<b>EMR EMT AEMT PARAMEDIC</b>
-------------------------------

The following conditions require patients to be transported by stretcher or stair chair. Other patients may be transported ambulatory unless their condition warrants stretcher use.

1. Pregnant greater than 20 weeks
2. Possible cardiac chest pain
3. Shortness of breath
4. Asthma/Chronic Obstructive Pulmonary Disease
5. Stroke
6. Patients requiring spinal immobilization
7. Penetrating trauma to the torso, neck, or head
8. Lower extremity, pelvis trauma
9. Low back trauma
10. Unconscious, unresponsive patients
11. Seizures within past hour or actively seizing
12. Generalized weakness
13. Patients unable to ambulate secondary to pain or weakness
14. Altered level of consciousness, except psychiatric patients
15. Psychiatric patients requiring restraint

MISCELLANEOUS

611 Terminally Ill Patients

EMR EMT AEMT PARAMEDIC

Standing Order

1. Maintain a calm environment and avoid performing measures beyond basic life support.
2. Elicit as much information from persons present who are familiar with the patient's condition as possible.
3. Obtain and document the name and telephone number of the patient's physician if possible.
4. Maintain BLS procedures and contact Medical Control as soon as possible. Provide full information on the patient's present condition, history, and name of the patient's physician and telephone number.
5. Medical Control will direct the management of the call 6. Accept DNR/POST forms (original or copy):
  - a. State approved forms
  - b. Signed order in patient's medical records: nursing home, hospice, or home care

**Note:** If DNR/POST for is used to withhold or terminate resuscitation efforts, a copy must be attached to the PCR.

## MISCELLANEOUS

### 612 “Excited Delirium” / Taser Use

#### Assessment

Changes in LOC Ongoing disorientation Agitation Hallucination Hyperthermia Seizure Chest pain or difficulty breathing Significant injury from fall or takedown
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

#### EMR

1. Oxygen and airway maintenance appropriate for patient’s condition
2. Supportive care
3. **EMR STOP**

#### EMT

4. Pulse oximetry
5. **EMT STOP**

#### AEMT

6. Glucose check
7. IV LR or NS, bolus (20 cc/kg)
8. Titrate Dextrose 50% PRN slowly until normal levels achieved. Try to avoid large swing in serum glucose levels.
9. **AEMT STOP**

#### PARAMEDIC

10. EKG Monitor, 12 Lead EKG if appropriate
11. Haldol 5-10 mg IM or Versed 5-10 mg IM
12. Versed 2-5 mg IVP if generalized seizure activity

#### NOTES:

- Remember to always stay safe when dealing with excited delirium patients and have enough personnel available to safely and adequately subdue the patient. Do not attempt to further agitate the patient by attempting to start an IV if it cannot be done safely. Utilize IM medication and adequate personnel to provide care.
- All persons subjected to use of the device should be medically evaluated and monitored regularly.

### MISCELLANEOUS

- Darts should be treated as biohazard, and not be removed in the field except by trained personnel.
- Darts to eyes, mouth, face, neck and genitals or near indwelling medical devices or lines should not be removed in the field.

## MISCELLANEOUS

### 613 Neonatal Resuscitation

#### Assessment

Newborn with respiratory or circulatory distress

#### EMR

1. Dry and place in face up head down position
2. Keep infant level with mother until cord is clamped
3. Suction airway, if obvious obstruction to spontaneous breathing or requiring or requiring positive pressure ventilation
4. Respirations
  - A. If spontaneous
    - I. Wait 1-2 minutes then complete clamping cord and cut between clamps
    - II. Cover infant head
    - III. Wrap and keep warm
    - IV. Provide Oxygen
    - V. Transport without delay
  - B. If no respirations
    - I. Stimulate respirations: rub back, snap bottom of feet gently, if no change or respirations become depressed (<20 bpm)
    - II. Re-suction airway
    - III. High flow oxygen if no change ventilate with BVM at 30/min
    - IV. Wait 1-2 minutes then clamp cord and cut between clamps
    - V. Transport Immediately
5. Pulse
  - A. If pulse rate is less than 60 perform CPR at rate of 120 compressions/min, transport
6. **EMR STOP**

#### EMT

7. Pulse oximetry
8. **EMT STOP**

#### AEMT

9. INT or IV NS, if hypotensive bolus 20 cc/kg
10. The dose of epinephrine is 0.01 mg/kg IV/IO (0.1 cc/kg of 1:10,000) given q 3-5 minutes, and repeat until heart rate is above 60/minute. Refer to the length based tape to confirm dosage.
11. If pulse rate is >60 keep warm, ventilate with BVM if necessary, transport
12. **AEMT STOP**

MISCELLANEOUS

**PARAMEDIC**

13. EKG Monitor, 12 Lead EKG, if appropriate

## MISCELLANEOUS

### Capnography

#### PARAMEDIC

##### Indications:

- Capnography shall be used as soon as possible in conjunction with any airway management adjunct, including endotracheal, cricothyrotomy, Blind Insertion Airway Device (BIAD) or BVM
- Capnography is recommended to be used on all patients treated with CPAP, Magnesium, and/or Epinephrine for respiratory distress.

##### Procedure:

1. Attach capnography sensor to the BIAD, endotracheal tube, or oxygen delivery device.
2. Note CO<sub>2</sub> level and wave form changes. These will be documented on each respiratory failure, cardiac arrest, or respiratory distress patient.
3. Capnography shall remain in place with the airway and be monitored throughout the prehospital care and transport.
4. Any loss of CO<sub>2</sub> detection or waveform indicates an airway problem and should be documented.
5. Capnography should be monitored as procedures are performed to verify or correct the airway problem.
6. Document the procedure and results on/with the Patient Care Report (PCR/EPCR).
7. In all patients with a pulse, an ETCO<sub>2</sub>>20 is anticipated. In the post-resuscitation patient, no effort should be made to lower ETCO<sub>2</sub> by modification of the ventilatory rate. Further, in post-resuscitation patients without evidence of ongoing, severe bronchospasm, ventilatory rate should never be <6 breaths per minute.

In the pulseless patient, and ETCO<sub>2</sub> waveform with an ETCO<sub>2</sub> value >10 may be utilized to confirm the adequacy of an airway to include BVM and advanced devices with SpO<sub>2</sub> will not register.

MISCELLANEOUS

PROCEDURE

Chest Decompression

**PARAMEDIC**

1. Cleanse skin on affected side using aseptic technique
2. Using a 14 or 16 gauge 3 ½” angiocath, insert between the 2<sup>nd</sup>/3<sup>rd</sup> mid clavicular or 4<sup>th</sup>/5<sup>th</sup> mid-axillary spaces
3. Advance needle until “pop” is felt while the needle is entering the pleural space
4. Advance catheter until hub contacts skin
5. Cover catheter hub with Asherman Chest Seal (ensure one way valve effect)
6. Reassess patient for breath sound changes
7. If signs of tension reoccur check chest seal, consider repeating chest decompression per above steps
8. Contact Medical Control
9. Transport

*Use the same procedure for pediatric patients: use 18 or 20 gauge angiocath*

**PROCEDURE**

**Surgical Airway Procedure**

**PARAMEDIC**

**Three Step Cricothyrotomy**

**Equipment Needed:**

6.0 cuffed Endotracheal Tube

# 11 Scalpel

Bougie

**Process:**

**Step 1:**

**Skin Incision:** Quickly cleanse the neck, and grasp the larynx with the non-dominant hand. Use the index finger of the non-dominant hand to identify the thyroid cartilage, cricothyroid membrane, and cricoid ring. Once the underlying structures have been identified, use the dominant hand to make a vertical incision over the cricothyroid membrane. Place the non-dominant index finger into the vertical incision and move it side to side to clearly feel the cricothyroid membrane.

**Step 2:**

**Incision of Cricothyroid Membrane:** Remove the non-dominant index finger from the cricothyroid membrane, and make a 5-mm horizontal incision through the cricothyroid membrane. Watch the depth of incision to avoid injury to the underlying esophagus. Place the elastic bougie into the defect, and advance it until resistance is appreciated. This indicates entry into the right main stem bronchus.

**Step 3:**

**Endotracheal Tube Placement:** Advance the preselected cuffed endotracheal tube over the elastic bougie up to the cricothyroid membrane. Ensure that the bevel of the endotracheal tube is lined up with the horizontal incision of the cricothyroid membrane before advancing further. Apply gentle pressure while advancing the endotracheal tube through the divided cricothyroid membrane. As the bevel of the endotracheal tube is passing through the membrane, it will push the cricothyroid membrane laterally. This will open the defect, allowing placement of the larger endotracheal tube. Once the endotracheal tube cuff has entered the trachea, stop advancing. Remove the elastic bougie, and inflate the endotracheal tube cuff. Verify placement as with normal intubation.

## Continuous Positive Airway Pressure (CPAP)

### PARAMEDIC

Continuous Positive Airway Pressure has been shown to rapidly improve vital signs, gas exchange, reduce the work of breathing, decrease the sense of dyspnea, and decrease the need for endotracheal intubation in patients who suffer from shortness of breath from asthma, COPD, pulmonary edema, CO poisoning, Near Drowning, CHF, and pneumonia. In patients with CHF, CPAP improves hemodynamics by reducing left ventricular preload and afterload.

#### Indications

- Any patient who is respiratory distress for reasons other than trauma or pneumothorax, and;
- Is awake and able to follow commands
- Is over 12 years old and the CPAP mask fits appropriately
- Has the ability to maintain an open airway
- Has a systolic blood pressure above 90 mmHg
- Uses accessory muscles during respirations
- Shows signs and symptoms consistent with asthma, COPD, pulmonary edema, CHF or pneumonia

**AND** who exhibit **two or more** of the following:

- A respiratory rate greater than 25 breaths per minute
- Pulse Oximetry of less than 94% at any time
- Use of accessory muscles during respirations

#### Contraindications

- Patient is in respiratory arrest/apneic
- Patient is suspected of having a pneumothorax or has suffered trauma to the chest
- Patient has a tracheostomy
- Patient is actively vomiting or has upper GI bleeding
- Patient has decreased cardiac output, obtundation and questionable ability to protect airway (e.g. Stroke, etc), penetrating chest trauma, gastric distention, severe facial injury, uncontrolled vomiting, and hypotension secondary to hypovolemia

#### Precautions

Use care if patient:

- Has impaired mental status and is not able to cooperate with the procedure
- Has failed at non-invasive ventilation
- Has active upper GI bleeding or history
- Complains of nausea or vomiting
- Has inadequate respiratory effort
- Has excessive secretions
- Has a facial deformity that prevents the use of CPAP

**CONTINUED ON NEXT PAGE**

## Procedure

Explain the procedure to the patient

1. Connect O<sub>2</sub> tubing nipple to gas source
2. Place the face mask securely to the patient's face using head harness
3. With nebulizer in the OFF position slowly increase gas flow to 6 or 8 LPM. Check face mask fit to patient and device connections for leaks.
4. Adjust the flow meter until desired pressure is obtained. **Maximum benefit is usually achieved at about 7.5 mm H<sub>2</sub>O. Higher pressures result in more side effects with minimal improvements in benefits.** Flow of 12-14 LPM is required to reach CPAP pressure of 8.5-10 cm H<sub>2</sub>O
5. Do not exceed 33 LPM
6. Patient SaO<sub>2</sub> should be monitored using a pulse oximeter.
7. To activate nebulizer rotate knob to the ON position.
8. If necessary, readjust flow meter to obtain desired CPAP pressure. Up to 25 LPM may be required.
9. Consider Ondansetron (Zofran) 2 - 4 mg IV (*peds 0.15 mg/kg IV*)

## Measuring Pressure

- Pressure relief limits maximum CPAP pressure to 25 cm H<sub>2</sub>O @ 25 LPM
- Do not exceed pressure limit of manometer (25 cm H<sub>2</sub>O)
- Manometer accuracy ± 3 cm H<sub>2</sub>O up to 15 cm H<sub>2</sub>O and ± 5 cm H<sub>2</sub>O over 15 cm H<sub>2</sub>O

## Notes:

- In the event of undesirable flow from oxygen source, simply remove the device and place on supplemental oxygen.
- Use of the Flow-Safe with non-back pressure compensated flow devices may affect input gas liter flow. Always verify delivered CPAP pressure on a manometer.
- Activation or deactivation of nebulizer may affect the delivered CPAP pressure. Always verify delivered CPAP pressure with a monometer.
- Flow meters capable of delivering up to 25 LPM may be required to operate both CPAP and Nebulizer simultaneously.
- Use of nebulizer other than the one provided may affect performance.
- Do not remove CPAP until hospital therapy is ready to be placed on the patient.
- Watch the patient for gastric distention that can result in vomiting
- Procedure may be performed on patients with a Do Not Resuscitate order
- Due to the changes in preload and afterload of the heart during CPAP therapy, a complete set of vital signs must be obtained every 5 minutes.

## Delayed Off Loading of Stable Non-Emergent Patients in the ED

EMS is currently facing an increasing frequency of patient turnover being delayed in the Emergency Department due to delays in acknowledgement, assessment, and placement in the ED. These delays negatively impact the ability of EMS to maintain response capability and provide emergency response in a timely manner. This protocol provides a method to off-load non-emergent patients and return to service in a timely manner.

AEMT
PARAMEDIC

Eligible patients (patients must meet ALL of the following criteria:

- Greater than 16 years old or less than 65 years old
- Stable vital signs
- Non-Emergent complaint
- Patient can walk and talk
- Patient has had neither medications nor significant interventions by EMS (minor bandaging, splinting, without Nausea/Vomiting concerns)

### Procedure

1. Ambulance arrives in ED and notifies ED nursing staff of patient.
2. If the ED Nursing Staff has not accepted report and made efforts to offload the patient from the EMS stretcher within 30 minutes of arrival, contact EMS Supervisor.
3. EMS Supervisor again requests ED Nursing Staff to offload the EMS stretcher. If no progress is made within 15 minutes of the Supervisor's engagement, and the patient meets all the criteria above, perform the following:
  - a. Ensure the patient's condition is unchanged
  - b. Walk the patient to the triage waiting area
  - c. Document all contacts with ED personnel, and record names of Charge and Triage nurses
  - d. Denote method of patient care transfer on PCR/EPCR
  - e. Complete an abbreviated, hand written EMS run report to include patient demographics, complaint, vital signs and pertinent history and ensure hospital is aware of patient's presence in the waiting room
  - f. Complete standard EPCR run report
  - g. Return to service

In the event that a patient does not agree to be placed in the waiting room the patient has the right to refuse offload.

## Endotracheal Tube Introducer (Bougie)

### PARAMEDIC

#### Indications:

- Patients meet clinical indications for oral intubation (appropriate to use with any attempt)
- Predictable difficult intubation

#### Contraindications:

- Introducer larger than ETT internal diameter

#### Procedure:

1. Prepare, position and oxygenate the patient with 100% Oxygen
2. Select proper ET tube without stylet, test cuff and prepare suction
3. Lubricate the distal end and cuff of the endotracheal tube and the distal ½ of the Endotracheal Tube Introducer (Bougie) (Note: failure to lubricate the Bougie and the ETT may result in being unable to pass the ETT)
4. Using laryngoscopic techniques, visualize the vocal cords if possible using Sellick's/ BURP as needed
5. Introduce the Bougie with curved tip anteriorly and visualize the tip passing the vocal cords or above the arytenoids if the cords cannot be visualized
6. Once inserted, gently advance the Bougie until you meet resistance or "hold-up" (if you do not meet resistance you have a probable esophageal intubation and insertion should be reattempted or the failed airway protocol implemented as indicated)
7. Withdraw the Bougie only to a depth sufficient to allow loading of the ETT while maintaining proximal control of the Bougie
8. Gently advance the Bougie and loaded ET Tube until you have hold-up again, thereby assuring tracheal placement and minimizing the risk of accidental displacement of the Bougie
9. While maintaining a firm grasp on the proximal Bougie, introduce the ET Tube over the Bougie passing the tube to its appropriate length
10. If you are unable to advance the ETT into the trachea and the Bougie and ETT are adequately lubricated, withdraw the ETT slightly and rotate the ETT 90° COUNTER CLOCKWISE to turn the bevel of the ETT posteriorly. If this technique fails to facilitate passing of the ETT you may attempt direct laryngoscopy while advancing the ETT (this will require an assistant to maintain the position of the Bougie and if so desired advance the ETT)
11. Once the ETT is correctly placed, hold the ET Tube securely and remove the Bougie
12. Confirm tracheal placement, inflate the cuff with 3-10 cc of air, auscultate for equal breath sounds and reposition accordingly

When final position is determined, secure the ET Tube, reassess breath sounds, apply end tidal CO<sub>2</sub> monitor, and record the monitor readings to assure continued tracheal intubation

## External Transcutaneous Cardiac Pacing

### PARAMEDIC

Noninvasive cardiac pacing, also referred to as external or transcutaneous pacing, involves the temporary application of externally applied electrodes to deliver an adjustable electrical impulse directly across an intact chest wall for the purpose of rhythmically stimulating the myocardium to increase the mechanical heart rate.

#### Indications:

- It is indicated for the treatment of hemo-dynamically compromised patients in settings where cardiac output is compromised due either to the complete failure of cardiac rhythm or to an insufficient rate of the patient's intrinsic pacemaker.
- Bradycardia with a systolic BP of less than 80 mmHg with shock-like signs or symptoms.
- Patients who experience provider-witnessed cardiopulmonary arrest and who present with asystole, or patients whose EKG converts to asystole while the EKG is being monitored.
- Prompt application of the transcutaneous cardiac pacemaker is appropriate prior to the administration of epinephrine and atropine when a patient converts to asystole as a primary rhythm during EKG monitoring by an EMT-P.
- Pediatric patients (40 kg or less) with profound symptomatic bradycardia unresponsive to optimal airway management, oxygenation, epinephrine, and atropine.

**NOTE:** Medical consultation is required for pacing pediatric patients.

#### Contraindications:

- Non-witnessed cardiopulmonary arrest with asystole
- Patients not meeting blood pressure criteria

#### Technique:

Start at a pacemaker heart rate of 70 beats per minute and the milliamperes (m.a.) as low as possible. Gradually increase m.a. until palpable pulse confirmed capture or 200 m.a.

#### Potential Adverse Effects/Complications:

Patients may experience mild to moderate discomfort. If patient is conscious and has adequate blood pressure consider:

- **Morphine** 2-10 mg IV/IO or **Fentanyl** 25-50 mcg IV/IO
- **Versed** 2-4 mg IV/IO

Musculoskeletal twitching in the upper torso may occur during cardiac pacing.

#### Precautions:

When properly applied, chest compressions can be performed directly over the insulated electrodes while the pacer is operating.

**DO NOT USE EXTERNAL CARDIAC PACING ON A HYPOTHERMIC PATIENT.**

## EZ-IO

AEMT

PARAMEDIC

### Indications:

1. Intravenous fluid or medications needed AND
2. Peripheral IV cannot be established in two attempts or 90 seconds AND the patient exhibits one or more of the following:
  - a. Altered mental status (GCS of 8 or less)
  - b. Respiratory compromise (SaO<sub>2</sub> of 80% or less following appropriate oxygen therapy, and/or respiratory rate <10 or >40/min)
  - c. Hemodynamically unstable (Systolic BP <90)
3. IV Access is preferred, however, IO may be considered prior to peripheral IV attempts in the following situations:
  - a. Cardiac Arrest (Medical or Trauma)
  - b. Profound hypovolemia with altered mental status

### Contraindications:

1. Fracture of the tibia or femur (for tibia insertion) - Consider alternate tibia
2. Fracture of the humerus (for humeral head insertion) - Consider alternate humerus
3. Previous orthopedic procedures (ex.: IO within previous 24 hrs, knee replacement, shoulder replacement)
4. Infection at the insertion site
5. Significant edema
6. Excessive tissue at insertion site
7. Inability to locate landmarks

### Considerations:

1. Flow rates: Due to the anatomy of the IO space you will note flow rates to be slower than those achieved with IV access
  - a. Ensure the administration of 10 ml rapid bolus with syringe
  - b. Use a pressure bag or pump for fluid challenge
2. Pain: Insertion of the IO device in conscious patients causes mild to moderate discomfort and is usually no more painful than a large bore IV. However, fluid infusion into the IO space is very painful and the following measures should be taken for conscious patients:
  - a. Prior to IO bolus or flush on a conscious adult patient, SLOWLY administer 20-50 mg of 2% lidocaine.
  - b. **Prior to IO bolus or flush on a conscious pediatric patient, SLOWLY administer 0.5 mg/kg 2% lidocaine.**

CONTINUED ON NEXT PAGE

# Roane County OES, EMS Division

---

## Adult patient:

- Defined as a patient weighing 40 kg or greater
- The adult (blue cap) needle set shall be used for adult patients

## Primary Insertion Site: Tibial Plateau

If IO access is warranted, the tibia shall be the insertion site of choice if possible

**Note:** In the cardiac arrest patient the humeral head should be the primary insertion site

## Alternate Insertion Site: Humeral Head or Medial Malleolus (adult patient only)

If IO access is not available via the tibial insertion site due to contraindications or inability to access the site due to patient entrapment and vascular access is imperative, the IO may be placed in the humeral head.

## Notes:

- In the cardiac arrest patient the humeral head should be the primary insertion site
- Do not attempt insertion medial to the Intertubercular Groove or the Lesser Tubercle

## Pediatric Patient:

- Defined as a patient weighing 3-39 kg
- The pediatric needles set (pink cap) shall be used for pediatric patients
- Use the length based assessment tape to determine pediatric weight
- The only approved site for pediatric IO insertion is the tibial plateau

## Standing Order:

The EZ-IO may be used if the indications are met and no contraindications exist.

## Precautions:

- The EZ-IO is not intended for prophylactic use
- The EZ-IO infusion system requires specific training prior to use
- Proper identification of the insertion site is crucial.

## Landmarks: Tibial Plateau

There are three important anatomical landmarks - the Patella, the Tibial Tuberosity (if present) and the Flat Aspect of the Medial Malleolus.

- **Important: The tibial tuberosity is often difficult or impossible to palpate on very young patients!**
- The traditional approach for IO insertions in small patients where the tibial tuberosity cannot be palpated is to identify the insertion site **“TWO FINGER WIDTHS BELOW THE PATELLA and then medial along the flat aspect of the TIBIA”**.
- The traditional IO insertion in slightly larger patients where the tuberosity can be appreciated generally suggests **“One finger width distal to the tibial tuberosity along the flat aspect of the medial tibia.”**
- The EZ-IO should be inserted two finger widths below the patella (kneecap) and one finger medial (toward the inside) to the tibial tuberosity.
- **For the morbidly obese patient:**
  - Consider rotating the foot to the mid-line position (foot straight up and down).
  - With the knee slightly flexed, lift the foot off the surface allowing the lower leg to “hang” dependent.

## Roane County OES, EMS Division

---

- o This maneuver may improve your ability to visualize and access the tibial insertion site.
  - o **Landmarks: Humeral Head**
    - Place the patient in a supine position
    - Expose the shoulder and place the patient's arm against the patient's body.
    - Rest the elbow on the stretcher with the forearm on the abdomen. Palpate and identify the mid shaft humerus and continue palpating toward the humeral head. As you near the shoulder you will note a small protrusion. This is the base of the greater tubercle insertion site. With the opposite hand "pinch" the anterior and inferior aspects of the humeral head confirming the identification of the greater tubercle. This will ensure that you have identified the midline of the humerus itself. The insertion site is approximately two finger widths inferior to the coracoid process and the acromion.

### **Landmarks: Medial Malleolus**

- The insertion site is two finger widths proximal to the Medial Malleolus and positioned midline on the medial shaft

### **Procedure:**

#### Inserting the EZ-IO:

1. Determine that the EZ-IO is indicated
2. Ensure that no contraindications are present
3. Locate the proper insertion site
4. Clean the insertion site with alcohol
5. Prepare the EZ-IO driver and needle set
6. Stabilize the leg (or arm)
7. Position the driver at the insertion site with the needle at a 90° angle to the surface of the bone
8. Power the needle set through the skin until you feel the tip of the needle set encounter the bone. Apply firm steady pressure on the driver and power through the cortex of the bone. Stop when the needle flange touches the skin or a sudden resistance is felt. **Stop on the POP.** This indicates entry into the bone marrow cavity
9. Grasp the hub firmly with one hand and remove the driver from the needle set
10. While continuing to hold the hub firmly, rotate the stylet counter clockwise and remove it from the needle set. Dispose of the stylet properly in a sharps container
11. Confirm proper placement of the EZ-IO catheter tip:
  - a. The catheter stands straight up at a 90° angle and is firmly seated in the tibia
  - b. Blood is sometimes visible at the tip of the stylet
  - c. Aspiration of a small amount of marrow with a syringe
12. Attach a primed extension set to the hub and flush the IO space with 10 cc of Normal Saline. **NO FLUSH - NO FLOW**
13. If the patient is conscious, administer Lidocaine 2% 20-50 mg slowly **PRIOR** to the initial bolus
14. Initiate the infusion per standing orders. Use of a pressure infuser or blood pressure cuff is recommended to maintain adequate flow rates **15.**Apply the wrist band and a dressing

## Roane County OES, EMS Division

---

**NOTE:** With properly documented training and equipment AEMTs are authorized to place pediatric IOs. Pediatric IOs should be utilized in accordance with these protocols.

## PROCEDURE

### Intranasal Medication

AEMT

PARAMEDIC

Medication administration in a certain subgroup of patients can be a very difficult endeavor. For example, an actively seizing or medically restrained patient may make attempting to establish an IV almost impossible which can delay effective drug administration. Moreover, the paramedic or other member of the medical team may be more likely to suffer a needlestick injury while caring for these patients.

In order to improve prehospital care and to reduce the risks of accidental needle-stick, the use of Mucosal Atomizer Device (MAD) is authorized in certain patients. The MAD allows certain IV medications to be administered into the nose. The device creates a medication mist which lands on the mucosal surfaces and is absorbed directly into the bloodstream.

#### Indications:

Emergent need for medication administration and IV access unobtainable or presents a high risk of needle-stick injury due to patient condition

- Seizures/Behavioral control: Midazolam (Versed) may be given intranasally until IV access is available
- Altered Mental Status from Suspected Narcotic Overdose: Naloxone (Narcan) may be given intranasally until IV access is available

Medications administered via the IN route require a higher concentration of drug in a smaller volume of fluid than typically used in the IV route. In general, no more than 1 milliliter of volume can be administered during a single administration event.

#### Contraindications:

- Bleeding from the nose or excessive nasal discharge
- Mucosal destruction

#### Technique:

1. Draw proper dosage (see below)
2. Expel air from syringe
3. Attach the MAD device via LuerLock Device
4. *Briskly* compress the syringe plunger

#### Complications:

- *Gently* pushing the plunger will not result in atomization
- Fluid may escape from the nares
- IntraNasal Dosing is less effective than IV dosing (slower onset, incomplete absorption)
- Current patient use of nasal vasoconstrictors (Neosynephrine/Cocaine) will significantly reduce the effectiveness of IN medications. Absorption is delayed, peak drug level is reduced, and time of drug onset is delayed.

# Roane County OES, EMS Division

---

## Midazolam (Paramedic Only)

### Precautions:

1. Midazolam may cause hypoventilation and potential respiratory depression/arrest. Have equipment and help readily available to manage the airway when administering this medication.
2. If hypotension develops after the administration of Midazolam, administer a 20 ml/kg bolus of normal saline.

Patient Age (years)	Weight (kg)	IN Midazolam volume in ml (assuming 5 mg/ml concentration) Midazolam volume dose (mg)
Neonate	3	0.12 ml 0.6 mg
<1	6	0.24 ml 1.2 mg
1	10	0.40 ml 2.0 mg
2	14	0.56 ml 2.8 mg
3	16	0.64 ml 3.2 mg
4	18	0.72 ml 3.6 mg
5	20	0.80 ml 4.0 mg
6	22	0.88 ml 4.4 mg
7	24	0.96 ml 4.8 mg
8	26	1.04 ml 5.2 mg
9	28	1.12 ml 5.6 mg
10	30	1.20 ml 6.0 mg
11	32	1.28 ml 6.4 mg
12	34	1.36 ml 6.8 mg
Small Teenager	40	1.60 ml 8.0 mg
Adult or Full grown teenager	50 or more	2.0 ml 10.0 mg

### Naloxone (Paramedic and AEMT Only) Adult:

1. Naloxone 0.4 mg every 5 minutes until the respiratory rate improves and the patient can maintain a pulse oximetry reading of 96% OR until 2 mg has been given

### Pediatric:

- A. Naloxone 0.1 mg/kg (max single dose 0.4 mg) until the respiratory rate improves and the patient can maintain a pulse oximetry reading of 96% OR until 2 mg has been given

## PROCEDURE

### Rapid Sequence Intubation

## PARAMEDIC

### Assessment and Indications

- GCS  $\leq$  8
- Intact gag reflex with potential airway compromise
- Combativeness that threatens airway or spinal cord compromise
- Potential for airway compromise during transport (inhalation injury, unstable facial fractures, severe bleeding into oral cavity) ·
- Need for ventilatory assistance or airway protection secondary to other causes

### Contraindications

- Known allergy to agents
- Inability to secure the airway by other means if ET intubation is unsuccessful
- Pediatric patients under the age of 8

### Precautions

- Pregnancy
- Dehydration
- Respiratory disease
- Penetrating eye injury
- Fractures and crush injury
- Cardiac disease
- Neuromuscular disease
- Severe burns

- Glaucoma
- Multiple facial fractures or facial instability

**Equipment -**

- Gather and prepare all equipment prior to starting procedure
- All equipment should be age appropriate

**Prepare the patient**

- Provide inline stabilization of the head and neck in the trauma patient
- Consider removing the anterior portion of the cervical collar
- Position the patient for optimal visualization
- Establish at least 1 IV/IO access (2 preferred) and ensure patency
- Attach cardiac monitor and pulse oximeter
- Pre-oxygenate the patient with 100% Oxygen via Non-rebreather if the patient is breathing spontaneously AND
- Nasal cannula at 15 LPM for at least two minutes (use even if patient is apneic)
- Assist the patient with a BVM only if respirations are inadequate or patient is apneic

**Protocol**

1. Sedation
  - A. Versed 0.05-0.1 mg/kg

**OR**

- B. Etomidate 0.3mg/kg (pediatrics 20mg max)

**PROCEDURE**

**Rapid Sequence Intubation continued**

2. Paralytics
  - A. Norcuron (Vecuronium) 0.15 mg/kg

**OR**

- B. Rocuronium Bromide 1-1.5mg/kg
3. Orally intubate when patient is apneic and paralyzed.
4. If unable to intubate within 20 seconds, halt attempts, provide ventilatory assistance for 30-60 seconds and reattempt intubation. If unable to intubate on 2<sup>nd</sup> attempt after 20 seconds, STOP! Provide ventilatory assistance and switch to ALTERNATE AIRWAY!
5. If unable to pass the alternate airway or it is contraindicated, then move to the Surgical Airway Procedure.
6. If the patient is bradycardic or intubation induced bradycardia occurs, temporarily halt the intubation procedure. Hyperventilate with manual resuscitator and high flow oxygen. If bradycardia continues administer Atropine 0.5 mg IVP, If bradycardia continues give 0.5 mg Atropine IVP every 5 minutes until it resolves or a maximum dose of 3.0 mg has been reached. *Pediatric Atropine: 0.02mg/kg with a minimum single dose of 0.1 mg and a maximum single dose of 0.5 mg. Maximum dose of 1 mg for a Child and 3 mg for an Adolescent.*
7. Continued Sedation, Paralysis and Pain control
  1. Versed 0.01 - 0.05 mg/kg every 15 minutes as needed
  2. Norcuron 0.01 - 0.05 mg/kg PRN for continued paralysis

**OR**

  3. Rocuronium 0.5 - 0.75 mg/kg

4. Consider giving Fentanyl 0.5 - 1 mcg/kg as an adjunct to Versed and for pain control

## PROCEDURE

### Rapid Sequence Intubation continued

#### Verify correct ETT placement

1. Visualize vocal cords during ETT placement
2. Auscultate thorax and abdomen to determine if air entry is adequate and symmetrical to all lung fields and absent over the epigastrium
3. Observe for symmetric chest wall expansion with ventilation
4. **\*\*Apply ETCO2 detector and visualize capnography**
5. Secure ETT at appropriate CM mark at lips in accordance with ETT size

#### Documentation

- Indication for intubation
- Tube size
- Pre-oxygenation prior to intubation and oxygen saturation prior to and after procedure
- Classification and condition of airway: clear, emesis, blood, etc.
- Difficulty with the procedure, including number of attempts
- Depth of insertion and how the tube is secured
- Who performed the procedure
- Cricoid pressure
- Manual in-line immobilization of C-Spine for trauma patients
- Means by which patient was ventilated after intubation and oxygen delivered

## Roane County OES, EMS Division

---

- Cardiac rhythm
- Status of ETT after each movement of patient
- Status of tube at receiving facility; breath sounds, oxygen saturation, End tidal CO<sub>2</sub>, clinical improvement/stability
- Head and neck immobilized on all patients (medical and trauma) for tube security
- Complete required QA sheet

**NOTE: RSI may ONLY be performed by paramedics who have documented competency in this skill via written confirmation with the medical director.**

## PROCEDURE

### Tourniquet

**EMR EMT AEMT PARAMEDIC**

#### Indications:

- Life threatening arterial hemorrhage
- **Serious or life threatening** extremity hemorrhage and tactical consideration prevent the use of standard hemorrhage control techniques

#### Contraindications:

- Non - Extremity hemorrhage
- Proximal extremity location where tourniquet application is not practical

#### Procedure:

1. Place tourniquet proximal to wound
2. Tighten per manufacturer instructions until hemorrhage stops and/or distal pulses in affected extremity disappear
3. Secure tourniquet per manufacturer instructions
4. Note time of tourniquet application and communicate this to receiving care providers
5. Dress wounds per standard wound care protocol
6. If delayed or prolonged transport and tourniquet application time greater than 2 hours, contact medical control
7. Include Tourniquet in use in your report to the Trauma Center as soon as practical and in your documentation for the PCR

If bleeding persists consider applying second tourniquet or using a hemorrhage control clamp

# Roane County OES, EMS Division

---

## REFERENCE

### Trauma Score

<b>RESPIRATORY RATE</b>	10-24/min	4
	24-35/min	3
	>36/min	2
	1-9/min	1
	None	0
<b>RESPIRATORY EXPANSION</b>	Normal	1
	Retractive	0
<b>SYSTOLIC BLOOD PRESSURE</b>	>90 mmHg	4
	70-89 mmHg	3
	50-69 mmHg	2
	0-49 mmHg	1
	No Pulse	0
<b>CAPILLARY REFILL</b>	Normal	2
	Delayed	1
<b>Points to add to the RTS based on the GCS</b>		
14-15		5
11-13		4
8-12		3
5-7		2
3-4		1

# Roane County OES, EMS Division

---

## Glasgow Coma Scale

<b>Eye Opening</b>	
Spontaneous	4
Opening to voice	3
Response to pain	2
None	1
<b>Verbal</b>	
Oriented	5
Verbal confused	4
Inappropriate words	3
Incomprehensible sounds	2
None	1
<b>Motor</b>	
Obeys commands	6
Localizes pain	5
Withdraws (pain)	4
Flexion	3
Extension	2
None	1

# Roane County OES, EMS Division

---

## REFERENCE

### Common Medical Abbreviations

<b>a</b> = before <b>AED</b> = Automated External Defibrillator	bladder <b>Gm/g</b> = gram
<b>AOX3</b> = alert and oriented to person place and time	<b>gr.</b> = grain
<b>Abd</b> = Abdomen <b>Ab.</b> = Abortion <b>ac</b> = antecubital <b>AF</b> = atrial fibrillation	<b>GSW</b> = Gunshot Wound
<b>ARDS</b> = Adult Respiratory Distress Syndrome	<b>gtt.</b> = drop <b>GU</b> = genitourinary <b>GYN</b> = gynecologic <b>h/hr</b> = hour <b>H/A</b> = headache
<b>AT</b> = atrial tachycardia	<b>Hg</b> = mercury
<b>AV</b> = atrioventricular	<b>H&amp;P</b> = History and Physical
<b>b.i.d.</b> = twice a day	<b>Hx</b> = history
<b>BSA</b> = Body Surface Area	<b>ICP</b> = intracranial pressure
<b>BS</b> = Blood sugar and/or Breath Sounds <b>c</b> = with <b>CC</b> or <b>C/C</b> = Chief Complaint	<b>JVD</b> = jugular venous distension
<b>CHF</b> = Congestive Heart Failure <b>CNS</b> = Central Nervous System <b>c/o</b> = complains of <b>CO</b> = Carbon Monoxide	<b>KVO</b> = keep vein open
<b>CO<sub>2</sub></b> = Carbon Dioxide	<b>LAC</b> = laceration
<b>D/C</b> = discontinue	<b>LBBB</b> = left bundle branch block
<b>DM</b> = diabetes mellitus	<b>MAEW</b> = moves all extremities well <b>NaCl</b> = sodium chloride
<b>DTs</b> = delirium tremens	<b>NAD</b> = No apparent distress/no acute distress
<b>DVT</b> = deep venous thrombosis	<b>NPO</b> = Nothing by mouth
<b>Dx</b> = Diagnosis	<b>NKA</b> = No known allergies
<b>ECG/EKG</b> = electrocardiogram	<b>OD</b> = overdose
<b>EDC</b> = estimated date of confinement	<b>O.D.</b> = right eye
<b>EJ</b> = external jugular	<b>O.S.</b> = left eye
<b>ENT</b> = ear, nose, and throat	<b>PERL</b> = pupils equal and reactive to light
<b>ETOH</b> = alcohol by definition is any chemical compound containing the Hydroxyl group OH. ETOH is the abbreviation of Ethanol (grain alcohol) <b>fl</b> = fluid <b>fx</b> = fracture <b>GB</b> = gall	<b>PID</b> = pelvic inflammatory disease
	<b>p.o.</b> = by mouth <b>1°</b> = primary, first degree <b>PTA</b> = prior to arrival <b>pt.</b> = patient <b>q</b> = every
	<b>q.h.</b> = every hour
	<b>q.i.d.</b> = four times a day <b>RBBB</b> = right bundle branch block
	<b>R/O</b> = rule out
	<b>ROM</b> = range of motion <b>Rx</b> = take, treatment <b>s</b> = without
	<b>S/S</b> = signs and symptoms <b>TIA</b> = transient ischemic attack
	<b>t.i.d.</b> = three times a day <b>V.S.</b> = vital signs
	<b>y.o.</b> = years old

# Roane County OES, EMS Division

---

## REFERENCE

### Drug Infusion Admix Dosage Guidelines

#### Lidocaine:

2 gram medication/500 mL D<sub>5</sub>W = 4 mg/mL (always use 60 gtt. Set)

1 mg/min = 15 gtt/min

2 mg/min = 30 gtt/min

3 mg/min = 45 gtt/min

4 mg/min = 60 gtt/min

#### Magnesium Sulfate:

4 gram in 250 cc D<sub>5</sub>W (16 mg/ml) run at 30-60 gtt/min

#### Dopamine:

400 mg /250 cc D<sub>5</sub>W or 800 mg/500 cc D<sub>5</sub>W = 1600 ug/mL (Always use 60 gtt. Set)

50 kg patient - 110 lbs    70 kg patient - 154 lbs    100 kg patient - 220 lbs

2.5 ug/kg/min =  
5 gtt/min

2.5 ug/kg/min =  
7 gtt/min

2.5 ug/kg/min =  
10 gtt/min

5 ug/kg/min =  
12 gtt/min

5 ug/kg/min =  
13 gtt/min

5 ug/kg/min =  
19 gtt/min

10 ug/kg/min =  
19 gtt/min

10 ug/kg/min =  
27 gtt/min

10 ug/kg/min =  
38 gtt/min

20 ug/kg/min =  
38 gtt/min

20 ug/kg/min =  
53 gtt/min

20 ug/kg/min =  
75 gtt/min

# Roane County OES, EMS Division

---

## REFERENCE

### PEDIATRIC SHOCK/TRAUMA

#### Pediatric Points to Remember

1. An infant is less than one year old
2. A child is from one to eight years old
3. Remember that few pediatric arrests are primary cardiac events. Most stem from respiratory (airway) problems, dehydration/metabolic, or hypothermia. Ensure that a child that arrests or that is pending arrest is well oxygenated, well hydrated and warm.
4. Prognosis is extremely poor for a child that arrests
5. Treat children aggressively before they arrest
6. When in doubt contact Medical Control
7. The use of a length based assessment tape is **required** for all pediatric patients as a guide for medication and equipment sizes
8. Remember that with children the Intraosseous drug route is quick to establish and may be easier than gaining IV access
9. Children may be effectively ventilated using a BVM. This is the preferred method of ventilation in respiratory or cardiac arrest

**\*\*\*\*If in doubt always contact Medical Control\*\*\*\***

#### Pediatric Trauma Score

(14 years of age and under)

Component	+2 points	+1 point	-1 point
Size	Greater than 20 kg	10-20 kg	Less than 10 kg
Airway	Normal	Oral/Nasal Airway	Unmaintainable/Intubated
Systolic BP	Greater than 90 mmHg	50-90 mmHg	Less than 50 mmHg
CNS	Awake	Obtunded/LOC	Coma
Open Wound	None	Minor	Major/Penetrating
Skeletal	None	Closed Fractures	Open/ Multiple Fractures

Total Point Values from Physical Presentation of Injury  
Trauma Score \_\_\_\_\_ Sum of Points

**AUTHORIZATION FOR STANDING ORDERS**

Emergency Medical Services (EMS) Standing Orders and Protocols (revision completed May 2014) are hereby adopted. They are to be initiated by EMS personnel within their scope of licensure whenever a patient presents with injury or illness covered by the protocols. Where indicated to contact Medical Control, the EMS Provider should receive voice orders from Medical Control before proceeding. Other orders may be obtained from Medical Control when the situation is not covered by the protocols or as becomes necessary as deemed by the EMT or Paramedic.

**Effective Date of these SOPs:** \_\_\_\_\_

\_\_\_\_\_  
EMS Service Medical Director

\_\_\_\_\_  
Date